Vol. 94, Sect. 5, 1957

ECHINODERMA

COMPILED BY

A. M. CLARK, M.A.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE FOUR SHILLINGS
April, 1959

SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The following are the Sections and their prices, viz .:-

					1		s.	d.
1. Compreh	ensive Z	oology		***	***		3	0
2. Protozoa	b			•••	•••		10	0
3. Porifera	•••						2	0
4. Coelente	rata	•••					4	0
5. Echinode	erma	• • •					4	0
6. Vermes							10	0
7. Brachion	ooda						4	0
8. Bryozoa							2	6
9. Mollusca		• • •		• • •			10	0
10. Crustace	a			•••	•••	•••	6	6
11. Trilobita		•••	•••		•••		5	0
12. Arachnic	la			•••			10	0
*13. Insecta		• • •			•••		50	0
14. Protocho							2	0
15. Pisces			•••				8	6
16A. Amphil							6	0
16B. Reptilia				***		•••	17	6
1		• • •			•••	•••	10	0
18. Mammal			•••			•••	10	0
19. List of N		ban and	Sylven	nomo	•••	• • •	4	0
IV. MINU OI I	ew dent	era and	bubge	uera		***		U

^{*} Obtainable only from the Commonwealth Institute of Entomology, 56, Queen's Gate, London, S.W.7.

5. ECHINODERMA

COMPILED BY

A. M. CLARK, M.A.

CONTENTS

						PAGE
I.	TITLES			 	 	 1
II.	Subject	INDEX		 	 	 11
III.	Systemat	CIC INDEX	•••	 	 	 14

I.—TITLES.

Direction 72. Completion and in certain cases correction of entries relating to the names of genera of the Phyla Mollusca, Brachiopoda, Echinodermata and Chordata made on the Official List of Generic Names in Zoology by Rulings given in Opinions rendered in the period up to the end of 1936. Opin. int. Comm. zool. Nom. 1E: 161-192.

Direction 73. Addition to the Official List of Specific Names in Zoology (a) of the specific names of twenty-five species of the Phyla Mollusca, Brachiopoda, Echinodermata and Chordata, each of which is the type species of a genus, the name of which was placed on the Official List of Generic Names in Zoology in the period up to the end of 1936 and (b) of the specific names of three species belonging respectively to the Phyla Brachiopoda, Echinodermata and Chordata, each of which is currently treated by some or all specialists as a senior subjective synonym of name of such a species. Opin. int. Comm. zool. Nom. 1E: 193-214.

Afzelius, B. A. (1). Basophilic structures in the cytoplasm of the sea urchin egg. Proc. Stockh. Conf. on Electron Microscopy 1956: 147-149 1 fig.

© Zoological Society of London, 1959 Vol. 94 Afzelius, B. A. (2). The acrosomal reaction of the sea urchin spermatozoon Proc. Stockh. Conf. on Electron Microscopy 1956: 167–169 4 figs.

Afzelius, B. A. (3). Electron microscopy on the basophilic structures of the sea urchin egg. Z. Zellforsch. **45**: 660–675 15 figs.

Afzelius, B. A. (4). Electron microscopy on sea urchin gametes. Uppsala (Almqvist & Wiksells) 20 pp.

Agrell, I. (1). The cellular division during the early embryonic development of the sea urchin embryo as influenced by temperature changes. K. fysiogr. Sällsk. Lund. Forh. 27: 61–72 6 figs.

Agrell, I. (2). Partial fertilization of the sea urchin egg by the action of oestradiol. Ark. Zool. 10: 491-496 5 figs.

[Aliev, R. A.] [Some species of corals and sea-urchins of the lower Cretaceous deposits of the rivers Tagirdzhalchaya and Kusarchaya.] C.R. Acad. Sci. Azerb. SSR 13: 1169–1175 pl. [In Russian.]

Amsden, T. W. Catalog of fossils from the middle and upper Ordovician of Oklahoma. Circ. Okla. geol. Surv. 43: 1-41.

Ancellin, J. Observations sur la faune et les fonds de pêche de quelques secteurs de la manche et des mers nordiques. Rev. Trav. Inst. Pêches marit., 21: 449-484 7 figs.

Anderson, J. M. see Guthrie, M. J.

Arvy, L. Contribution à la connaissance des "corps bruns" des Holothuridae. C.R. Acad. Sci. Paris 245: 2543-2545.

Bäckström, S. Content and distribution of ascorbic acid in sea urchin embryos of different developmental trends. Exp. Cell Res. 13: 333-340 7 figs.

Balinsky, J. B. The Ophiuroidea of Inhaca Island. Ann. Natal Mus. 14: 1-33 4 pls., 7 figs. map.

Banhidi, Z. G. An attempt to separate large amounts of sea urchin blastulae developed from embryonal fragments. Ark. Zool. 10: 449–452 1 fig.

Banse, K. Über den Transport von meroplanktischen Larven aus dem Kattegat in die Kieler Bucht. Ber. dtsch. Komm. Meeresforsch. 14 1956: 147–164 5 figs.

F[Baranova, Z. I.] [Echinoderms of the Bering Sea.] Invest. Far-east Seas U.S.S.R. 4: 149-266 19 figs. [In Russian.]

Barrett, J. H. see Bassindale, R.

Basch, P. F. Observations on the retractor strands of the starfish stomach. Biol. Rev. City Coll. N.Y. 18 1956: 14-17 6 figs.

Bassindale, R. & Barrett, J. H. The Dale Fort marine fauna. Proc. Bristol Nats. Soc. 29: 227–328 3 figs. (Echinoderma on pp. 313–317).

Bataller, J. R. La paleontologia y Luis Mariano Vidal. Bol. Inst. geol. Esp. 67 1956: 65–114 14 pls. (Echinoderma on pp. 29–33).

Bernasconi, I. (1). Equinoideos y Asteroideos de la Colección del Instituto Oceanográfico de la Universidad de San Pablo. Primera contrabución. Bol. Inst. oceanogr. S. Paulo 6 1955: 51-91 7 pls. [English summary.]

Bernasconi, I. (2). Otra nueva especie de Asteroideo brasileño. Neotropica 3 (11): 33–34 1 fig. Bernhard, M. Die Kultur von Seeigellarven (Arbacia lixula L.) in künstlichem und natürlichem Meerwasser mit Hilfe von Ionenaustauschsubstanzen und Komplexbildnern. Pubbl. Staz. zool. Napoli 29: 80–95 1 fig. 1 pl.

Blacker, R. W. Benthic animals as indicators of hydrographic conditions and climatic change in Svalbard waters. Fish. Invest. Min. Agric. London (2) 20 (10): 1-49 30 figs.

Blaschko, H. & Hope, D. B. Observations on and distribution of amine oxidase in invertebrates. Arch. Biochem. & Biophys. 69: 10-15.

Bolst, A. L. & Whiteley, A. H. Studies of the metabolism of phosphorus in the development of the sea urchin, Strongylocentrotus purpuratus. Biol. Bull. Woods Hole, 112: 276-287 2 figs.

Boroughs, H. see Hsiao, S. C.

Brookbank, J. W. see Tyler, A.

Cailleux, A. see Chavan, A.

Caso, M. E. (1). El genero Clypeaster Lamarck 1801, en el Terciario de Mexico. Resumenes de los Trabajos Presentados XX Congresso Geologico Internacional, Mexico 1956: 112.

Caso, M. E. (2). El genero *Clypeaster* Lamarck 1801, en el terciario de Mexico. An. Inst. Biol. Univ. Mex. 27: 487–528 23 figs.

Caster, K. E. & Eaton, T. H. Microstructure of the plates in the carpoid echinoderm *Paranacystis*. J. Paleont. 30 1956: 611-614 1 pl. 2 figs.

Caster, K. E. see Gill, E. D.

Chang, F.-Y. & Woo. P.-L. [On the echinoderms of Dairen and its vicinity.] Acta zool. Sinica 6 1954: 123-146 11 figs. 2 pls. [Chinese with English summary on p. 145.]

Chavan, A. & Cailleux, A. Détermination pratique des fossiles. Paris, 385 pp. figs.

Cherbonnier, G. (1). Une Ophiure nouvelle de Méditerranée: Amphiura apicula nov. sp. Vie et Milieu 8 2: 200-204 3 figs.

Cherbonnier, G. (2). Note sur une Holothurie dendrochirote: *Thyonacta* sabanilaensis (Deichmann). Bull. Mus. Hist. nat. Paris 28: 537-540 2 figs. Cherbonnier, G. (3). Ophiures rares ou nouvelles des côtes de Sierra-Leone. Bull. Mus. Hist. nat. Paris 29: 163-171 3 figs.

Choubert, H. & Termier, G. Présence du genre *Mimocystites* Barrande dans 'Ordovicien du Maroc. Notes Serv. Min. Maroc 117 1953: 137-143 7 figs.

Christensen, A. M. The feeding behavior of the seastar *Evasterias* troschelii Stimpson. Limnol. & Oceanogr. 2:180–197 15 figs.

Coe, W. R. Fluctuations in populations of littoral marine invertebrates. J. mar. Res. 15 1956: 212-232 9 figs.

Cognetti, G. Autofecondazione in Asterina. Boll. Zool. 23 1956: 275–278 1 pl.

Cognetti, G. see Delavault, R.

Collignon, J. Préliminaires à une étude écologique de la Baie de Pointe-Noire. Répartition de quelques Echinodermes caractéristiques. Bull. Inst. centafr. N.S. Nos. 13–14: 29–37 2 pls. 1 fig.

Conkin, J. E. Stratigraphy of the New Providence Formation (Mississippian) in Jefferson and Bullitt Counties, Kentucky, and fauna of the Coral Ridge member. Bull. Amer. Paleont. 38: 105-157 4 pls. 2 figs.

Cooke, C. W. (1). Geology of Saipan, Mariana Islands. Part 3. Paleontology Chapter J. Echinoids. Prof. Pap. U.S. geol. Surv. 280-J: 361-364 1 pl.

Cooke, C. W. (2). Rhynobrissus cuneus, a new echinoid from North Carolina. Proc. U.S. nat. Mus. 107 3379: 9-12 1 pl.

Cotronei, G. & Urbani, E. Ricerche sull' oocite di *Antedon rosacea* Lamarck. Pubbl. Staz. zool. Napoli **29**: 15–21.

Cramer, H. R. Ordovician starfish from the Martinsburg Shale, Swatara Gap, Pennsylvania. J. Paleont. 31: 903-907 1 pl. 1 fig.

Cumano, A. B. Astropecten aurantiacus (L.). Naturalia 4 1954: 158 fig.

Dan, J. C. The acrosome reaction. Int. Rev. Cytol. 5 1956: 365-393 21 figs. Deane, H. W. see Karnovsky, M. L.

Deflandre-Rigaud, M. Sur l'invalidité du genre *Theelia* Schlumb, synonyme de *Chiridotites* Defl.-Rig. (sclérites d'Holothurides fossiles). Bull. Mus. Hist. nat. Paris **29**: 353–355.

Deichmann, E. The littoral holothurians of the Bahama Islands. Amer. Mus. Novit. No. 1821: 1–20 71 figs.

Delavault, R. & Cognetti, G. L'hermaphrodisme chez *Echinaster sepositus* Gray du Golfe de Naples. C.R. Acad. Sci. Paris 245: 2545–2547 1 fig.

Devriès, A. Nouvelles observations sur *Infraclypeus thalebensis* Gauthier, Galeropygidé-Algérien. Bull. Soc. Hist. nat. Afr. N. **48**: 148–158 9 figs.

Dieuzeide, R., & Roland, J. Opérations de draguages et de chalutages effectuées au large des Côtes Algériennes au cours des années 1954 et 1955. Bull. Sta. Aquic. Pêche. Castiglione. No. 8 1956 1957: 11–27 2 figs.

[Djakonov, A. M.] [The echinoderm fauna of the Malayan Archipelego and its connection with the past and present forms of the Pacific.] Trud. Leningr. Soc. Nat. 70 1950: 109–126 2 figs. [In Russian.]

Domantay, J. S. Some holothurians from Guam and vicinity. Nat. appl. Sci. Bull. **12** 1954: 336–357.

Donovan, D. T. The Jurassic and Cretaceous systems in East Greenland. Medd. Grønland 155 4: 1-214 4 pls. 25 figs.

Druckrey, H., Schmähl, D. & Kaiser, K. Versuche mit zellteilungshemmenden Giften an unbefruchteten Seeigeleiern. Naturwissenschaften 44: 430-431.

Durham, J. W. Notes on Echinoids. J. Paleont. 31: 625-631 1 pl. 2 figs.

Durham, J. W. & Melville, R. V. (1). A classification of echinoids. Resumenes de los Trabajos Presentados XX Congresso Geologico Internacional, Mexico, 1956: 113-114.

Durham, J. W. & Melville, R. V. (2). A classification of Echinoids. J. Paleont. 31: 242–272 9 figs.

Eaton, T. H., see Caster, K. E.

[Eltuishcheva, R. S.] [On a new family of Paleozoic sea lilies.] Annu. Soc. paléont. russe 16: 218-235 3 pls. 9 figs.

Emerson, W. K. see Hertlein, L. G.

Endean, R. (1). The Cuverian tubules of Holothuria leucospilota. Quart. J. mier. Sci. 98: 455-472 9 figs.

Endean, R. (2). The biogeography of Queensland's shallow-water echinoderm fauna (excluding Crinoidea), with a rearrangement of the faunistic provinces of tropical Australia. Aust. J. mar. & freshw. Res. 8: 233-273 5 figs.

Engel, H. [in] Burck, H. D. M. Pliocene and lower Pleistocene in a boring near Oosterhout. Echinoidea. Overdruck Med. Geol. Sticht. n.s. 7 1953: 36-37 1 fig.

Engel, H. & Meijer, M. Notes sur les echinides du tuffeau de Maastricht (Maestrichtien, Dumont. 1849) Lychnidius scrobiculatus (Goldfuss, 1829). Natuurh. Maandbl. 46 7-8: 88-94 l pl. 4 figs.

Ephremova, L. N. see Postnikova, I. E.

Epis, R. C. see Langenheim, R. L. Exline, H. see Frizzell, D. L.

[Faas, A. V.] in [Librovitch, L.] [Atlas of the leading forms of the fossil fauna of the U.S.S.R. Vol. 4 Lower Carboniferous.] [Echinoidea] Leningrad 1941 pp. 72-75 pl. X (pt.). [In Russian.]

Falke, H. Zur Geologie der Umgebung von Stromberg (Hunsrück). Notizbl. hess. Landesamt. Bodenforsch. 85: 75-113 1 pl. 7 figs.

Fautrez, J. & Fautrez-Firlefyn, N. Hyperperméabilisation de l'oeuf immature d'Asterias glacialis sous l'influence du bleu de toluidine. C.R. Soc. Biol. Paris 151: 617-618.

Firtion, F. Les éléments paléontologiques dévoniens du Val de Bruche. Ann. Univ. saraviensis. Sci. 6: 97-184 10 pls.

Flickinger, R. A. Evidence from sea urchin-sand dollar hybrid embryos for a nuclear control of alkaline phosphatase activity. Biol. Bull. Woods Hole 112: 21-27 1 fig.

Frings, H. & Frings, M. Notes on Invertebrates found at Salisbury cove, Maine in July 1952. Bull. Mt. Desert Isl. Biol. Lab. 4 1953: 41-42.

Frizzell, D. L. & Exline, H. Revision of the family Synaptitidae, fossil holothurian sclerites (Echinodermata Holothuroidea), Bol. Soc. geol. Perú 32: 97-119 6 figs.

Galloway, J. J. & Kaska, H. V. Genus Pentremites and its species. Mem. geol. Soc. Amer. 69: 1-104 13 pls. 5 figs.

Gautier-Michaz, M. Cinq echinodermes nouveaux ou peu connus pour la faune de France. Rec. Trav. St. mar. Endoume No. 21: 80-83.

Ghekker, E. L. & Ghekker, R. F.] On a new species of the genus Protocrinites Eichwald.] Annu. Soc. paléont. russe 16: 274-279 1 pl. [In Russian.]

Gideon, P. W., Menon, P. K. B., Rao, S. R. V. & Jose, K. V. On the marine fauna of Gulf of Kutch: A preliminary survey. J. Bombay nat. Hist. Soc. 54: 690-706 1 pl. 5 figs. map.

Giese, A. C. & others. Reproductive cycles in some West Coast Echinoderms. Anat. Rec. 128: 554-555.

Gill, E. D. & Caster, K. E. Australian carpoid echinoderms. Resumenes de los Trabajos Presentados, XX Congresso Geologico Internacional, Mexico 1956: 116.

Grabert, G. & Grabert, H. Encrinaster schmidti (Schöndorf). Ein Leitfossil aus der Herdorf-Gruppe (oberes Siegenium). Paläont. Z. 30 1956: 190-198 2 pls. 2 figs.

Gruner, H.-E. see Stresemann, E.

Gustafson, T. & Hörstadius, Changes in the determination of the sea urchin egg induced by Amino Acids. Pubbl. Staz. zool. Napoli 29: 407-424 17 figs.

Guthrie, M. J. & Anderson, J. M. General Zoology. New York (John Wiley & Sons), pp. xv + 708, 1 chart, text illust. (Echinoderma on pp. 486-509.)

Hagström, B. E. Studies on polyspermy in sea urchins. Ark. Zool. 10: 307-315 1 fig.

Hagström, B. E. see Lybing, S.

Hagström, B. E. see Perlmann, P.

Hagström, B. E. & Markman, B. Further studies on the fertilization of jelly-free sea urchin eggs. Acta zool., Stockh. 38: 219-222 2 figs.

Hampton, J. S. Some Holothurian spicules from the Upper Bathonian of the Dorset coast. Geol. Mag. 94: 507-510 12 figs.

Hancock, D. A. The feeding behaviour of the sea urchin *Psammechinus miliaris* (Gmelin) in the laboratory. Proc. zool. Soc. Lond. **129**: 255-262 l pl.

Harris, B. see Hodson, F.

Hayashi, R. (1). Contributions to the classification of the sea-stars of Japan. I. Spinulosa. J. Fac. Sci. Hokkaido Imp. Univ. (6) Zool. 7 1940: 107-204 7 pls.

Hayashi, R. (2). A note of a Japanese sea-star, Astropecten ludwigi De Loriol. J. Fac. Sci. Hokkaido Univ. (Zool.) 13: 37-45 1 pl. 3 figs.

Hedgpeth, J. W. and others. Treatise on marine ecology and paleoecology. Vol. I. Ecology. Mem. geol. Soc. Amer. 67 1: 1-1296 figs.

Heller, F. Ein fund von *Trichasteropsis weissmanni* Münst. aus dem Oberen Muschelkalk bei Hassfurt/Ufr. Geol. Bl. NO. Bayern 7: 45–47, fig.

Hertlein, L. G. Pliocene and Pleistocene fossils from the Southern portion of the Gulf of California. Bull. S. Calif. Acad. Sci. 56: 57-75 1 pl.

Hertlein, L. G. & Emerson, W. K. Additional notes on the Invertebrate fauna of Clipperton Island. Amer. Mus. Novit. No. 1859: 1-9.

Hodson, F., Harris, B. & Lawson, L. Holothurian spicules from the Oxford Clay of Redcliff, near Weymouth (Dorset). Geol. Mag. 93 1956: 336– 344 25 figs.

Hodson, F. see Bradley, P. C. S.

Holter, H. & Zeuthen, E. Dynamics of early echinoderm development, as observed by phase contrast microscopy and correlated with respiration measurements. Pubbl. Staz. zool. Napoli 29: 285–306 3 pls. 7 figs.

Hope, D. B. see Blaschko, H.

Hörstadius, S. (1). Die Determination der bilateralen Symmetrie im Seeigelei. Verh. deuts. zool. Ges. 1956 1957: 193–194.

Hörstadius, S. (2). On the regulation of bilateral symmetry in plutei with exchanged meridional halves and in giant plutei. J. Embryol. exp. Morph. 5: 60-73 10 fgs.

Hörstadius, S. see Gustafson, T.

Hsiao, S. C. & Boroughs, H. The uptake of radioactive calcium by sea urchin eggs. Proc. Hawaii. Acad. Sci. No. 32: 27.

Hultin, T. (1). Acid-soluble nucleotides in the early development of *Psammechinus miliaris*. Exp. Cell Res. 12:413-415 2 figs.

Hultin, T. (2). The metabolic utilization of C¹⁴-formate in sea urchin embryos. Exp. Cell Res. **12**: 518–525 3 figs.

Immers, J. Cytochemical studies of fertilization and first mitosis of the sea urchin egg. Exp. Cell Res. 12: 145–153 13 figs.

Ishida, J. & Yasumasu, I. Changes in protein specificity determined by protective enzyme test during embryonic development of the sea-urchin and the fresh-water fish. J. Fac. Sci. Tokyo Univ. (Zool.) 8:95–107 8 figs.

Ishihara, K. Release and activation of aldolase at fertilization in sea-urchin eggs. J. Fac. Sci. Tokyo Univ. (Zool.) 8: 71–93 10 figs.

Jeffrey, S. S. see Karnovsky, M. L.

Johnson, J. H. Geology of Saipan, Mariana Islands. Part 2. Petrology and Soils. Chapter C. Petrography of the limestones. Prof. Pap. U.S. geol. Surv. 280-C: 173-187 3 pls. (Echinoderma on p. 180 + pl. xxxi.)

Jörg, E. Ein Erstfund von Asteriacites Schlotheim (Cubichnia, Asterozoa) aus der burdigalen Meeresmolasse des Bodenseegebietes. Beitr. naturk. Forsch. Karlsruhe 16: 34–36.

Jose, K. V. see Gideon, P. W.

Jux, U. Zwei neue Seesterne aus der Kreide von Abu Roâsch (Forbesiaster gaensis und Recurvaster lorioli) und das fazielle Bild des ägyptischen Santons. Paläont. Z. 31: 186–197 1 pl. 2 figs.

Kaiser, K. see Druckrey, H.

Karnovsky, M. L., Jeffrey, S. S., Thompson, M. S. & Deane, H. W. A chemical and histochemical studies of the lipides of the pyloric cecum of the starfish, Asterias forbesi. J. biophys. & biochem. Cytol. 1 1955: 173-182 1 pl. 2 figs. (Not seen by recorder.)

Kaska, H. V. see Galloway, J. J.

Kawaguti, S. & Yamasu, T. Carotenoid pigment in the test of cakeurchin, *Peronella japonica*. Biol. J. Okayama Univ. 3: 150–158 8 figs.

Kier, P. M. (1). Separation of interambulacral columns from the apical system in the echinoidea. J. Paleont. 30 1956: 971-974 3 figs.

Kier, P. M. (2). A new genus of echinoid from the Palaeozoic of Ireland. Geol. Mag. 93 1956: 15-17 1 pl. 2 figs.

Kier, P. M. (3). A new Upper Carboniferous Echinoid from Texas. Geol. Mag. 94: 326-328 1 fig.

Kier, P. M. (4). Tertiary Echinoidea from British Somaliland. J. Paleont. 31:839-902 5 pls. 20 figs.

Kinoshita, S. Changes in the phospholipid content of the starfish spermatozoa during their histidine induced movement. J. Fac. Sci. Tokyo Univ. (Zool.) 8: 41-49.

Kobayashi, N. & Nakamura, K. (1). [The cleavage inhibition & abnormal cleavage of sea-urchin eggs induced by Demecolcin I.] Zool. Mag. Tokyo 66: 258–262 4 figs. [Japanese with English summary.]

Kobayashi, N., & Nakamura, K. (2). [The cleavage inhibition and abnormal cleavage of sea-urchin eggs induced by demecolcin. II.] Zool. Mag., Tokyo. 66: 377–383 3 figs. [Japanese with English summary.]

Koenig, J. W. see Strimple, H. L.

Kongiel, R. Remarques sur les échinides suprajurassiques de Czarno-glow et de Swietoszewo en poméranie occidentale. Bull. Serv. géol. Pologne 105:5-747 pls. [In Polish, Russian and French.]

Krahl, M. E. Oxidative pathways for glucose in eggs of the sea urchin. Biochim. biophys. Acta 20 1956: 27–32.

Kriszat, G. & Runnström, J. Effects of dinitrophenols on the physical properties, developmental capacity and differentiation of sea urchin egg. Ark. Zool. 10: 595–606 3 figs.

Kriszat, G. see Runnström, J.

Kubo, K. Studies on the systematic serology in Sea-Stars I. J. Fac. Sci. Hokkaido Univ. (Zool.) 13: 67-71 3 figs. Kupka, E. Die Abhängigkeit der Kerngrössenvariation bei Seeigeln von verschiedenen physiologischen Bedingungen. Naturwissenschaften 44:122.

Ladd, H. S., and others. Treatise on marine ecology and paleoecology. Vol. 2. Paleoecology. Mem. geol. Soc. Amer. 67 2: 1-1077 figs.

Lallier, R. (1). Recherches sur l'animalisation de l'Oeuf de l'Oursin Paracentrotus lividus par les dérivés polysulfoniques. Pubbl. Staz. zool. Napoli 30: 185–209 11 figs.

Lallier, R. (2). Perchlorate et détermination embryonnaire chez les Echinodermes. C. R. Soc. Biol. Paris 151: 471-474.

Lallier, R. (3). Animalisation de l'oeuf de l'oursin *Paracentrotus lividus* par la polyanétholsulfonate de sodium (liquoide). C.R. Soc. Biol. Paris **151**: 638-641.

Lallier, R. (4). Colorants acides et détermination embryonnaire chez les echinodermes. Experientia 13: 362–364 figs.

Lallier, R. (5). Effets de substances sulfhydrilées sur la détermination embryonnaire de l'oeuf de l'oursin Paracentrotus lividus. C.R. Acad. Sci. Paris 246: 2810–2812.

Lallier, R. (6). Relations entre la structure de dérivés sulfoniques et leur action sur la détermination embryonnaire de l'oeuf de l'oursin *Paracentrotus lividus*. C. R. Acad. Sci. Paris 244: 3176–3178.

Langenheim, R. L. & Epis, R. C. Holothurian sclerites from the Mississippian Escabrosa limestone, Arizona. Micropaleontology 3: 165-170 1 pl. 1 fig.

Lawson, L. see Hodson, F.

Lehmann, W. M. Die Asterozoen in den Dachschiefern des rheinischen Unterdevons. Abh. hess. Landesamt. Bodenforsch. 21: 1-160 55 pls. 31 figs.

Liashenko, A. I. see Postnikova, I. E.

Lybing, S. & Hagström, B. E. Isolation of a fertilization-promoting factor from egg water of *Psammechinus miliaris*. Exp. Cell Res. 13: 60-68 3 figs.

Madsen, F. J. (1). Reports of the Lund University Chile Expedition 1948-49. 24. Asteroidea, with a survey of the Asteroidea of the Chilean Shelf. Acta. Univ. lund. Andra Afd. N.S. 52 2:1-53 6 pls.

Madsen, F. J. (2). On Walcott's supposed Cambrian Holothurians. J. Paleont. 31: 281-282.

Madsen, F. J. (3). On a new species of *Meoma*, an don a few other echinoids from tropical West Africa. Bull. Inst. franc. Afr. noire 19A: 474-481 3 figs.

Mancuso, V. La citocromo-ossidasi nell'uovo in sviluppo di *Sphaerechinus* granularis. Ric. sci. 27: 3049-3051. [Italian with French, English & German summaries.]

Marine Biological Association of the U.K. Plymouth Marine Fauna, Edit. 3, Plymouth (Mar. Biol. Assoc.) 457 pp. 4 charts (Echinoderma on pp. 358–369).

Markman, B. Studies on nuclear activity in the differentiation of the sea urchin embryo. Exp. Cell Res. 12: 424–426.

Markman, B. see Hagström, B. E.

Marshak, A. & Marshak, C. Thymine in the acid-soluble fraction of *Arbacia* eggs. J. biophys. & biochem. Cytol. 1 1955: 47-58 3 pls. (Not seen by recorder.)

Marsland, D. Protoplasmic contractility in relation to gel structure: Temperature pressure experiments on cytokinesis and amoeboid movement. Int. Rev. Cytol. 5 1956: 199–227 18 figs.

Meijer, M. see Engel, H.

Melville, R. V. see Durham, J. W.

Menon, P. K. B. see Gideon, P. W.

Messina, L. Prime osservazioni sulla natura chimica del liquido periviscerale di *Arbacia lixula*. Pubbl. Staz. Zool. Napoli 30: 127–131 1 fig.

Miller, H. W., Sternberg, G. F. & Walker, M. V. *Uintacrinus* localities in the Niobrara formation of Kansas. Trans. Kansas Acad. Sci. 60: 163-166 2 figs.

Millonig, G. I corpi vitellini nell' uovo di Arbacia lixula, osservati al microscopio elettronico. Monit. Zool. Ital., Suppl. 65: 354–356. Millott, N. (1). Naphthaquinone pigment in the tropical sea urchin Diadema antillarum Philippi. Proc. zool. Soc. Lond. 129: 263–272 l pl. 1 fig.

Millott, N. (2). Animal photosensitivity, with special reference to eyeless forms. Endeavour 16 (61): 19–28 21 figs.

Millott, N. see Vevers, H. G.

Millott, N. & Yoshida, M. The spectral sensitivity of the echinoid *Diadema* antillarum Philippi. J. exp. Biol. 34: 394-401 4 figs.

Mirigliano, G. Scutella montagnai n.sp. (Echinoidea, Clypeastroidea). Boll. Zool. 24: 9–15 1 pl.

Mitchison, J. M. Cell membranes and cell division. Symposia Soc. exp. Biol. No. 6 1952: 105–127 1 pl. 4 figs.

Miyauchi, Y. see Okada, K.

Mohri, H. (1). Fatty acid oxidation in sea-urchin spermatozoa. Annot. zool. jap. 30: 181–186 1 fig.

Mohri, H. (2). Endogenous substrates of respiration in sea-urchin spermatozoa. J. Fac. Sci. Tokyo Univ. (Zool.) 8: 51-63 3 figs.

Mohri, H. (3). Effect of malonate on oxygen uptake and dehydrogenase activity in sea-urchin spermatozoa. J. Fac. Sci. Tokyo Univ. (Zool.) 8:65–69 3 figs.

Monroy, A. (1). Ricerche sul significato fisiologico della capsula dell'uovo degli Echinodermi. R.C. Ist. sup. Sanit. 19 1956: 101–108 l pl.

Monroy, A. (2). Swelling properties of the mitochondria of unfertilized and newly fertilized sea urchin eggs. Experientia 13: 395–399 2 figs.

Monroy, A. see Nakano, E.

Moore, A. R. (1). Biparental inheritance in an interordinal cross of sea urchin and sand dollar. J. exp. Zool. 135: 75-79 2 pls.

Moore, A. R. (2). Morphogenetic modifications in the sea urchin by means of trypsin. Pubbl. Staz. zool. Napoli 29: 71–79 2 pls.

Morishita, A. (1). On some fossil echinoids from Kyusyu, Japan. Mem. Coll. Sci. Kyoto Univ. 23B 1956: 193-202 4 pls.

Morishita, A. (2). On some fossil echinoids of Japan. I. Mem. Coll. Sci. Kyoto Univ. 24B: 161-164 l pl.

Motomura, I. On the nature and localization of the third factor for the toughening of the fertilization membrane of the sea urchin egg. Sci. Rep. Töhoku Univ. (4) 23: 167–181 11 figs.

Mugard, H. L'infraciliature chez la larve d'oursin Paracentrotus lividus. Bull. Soc. Zool. France 82: 81-88 11 figs.

Nakamura, K. see Kobayashi, N.

Nakano, E. & Monroy, A. A method for the incorporation of radioactive isotopes in the sea urchin egg. Experientia 13: 416–417.

Nishibori, K. Studies on pigments of marine animals. III. Echinochrome A from the spines of sand-dollar, *Echina-rachnius mirabilis*. Bull. Jap. Soc. Sci. Fish. 22: 708–712 5 figs.

Okada, K. & Miyauchi, Y. Normal table of the early developmental stages on the sea-urchin, *Hemicentrotus pul-cherrimus*. J. Gakugei Tokushima Univ. Nat. Sci. 5 1954: 57-69 6 pls.

Panning, A. Bemerkungen über die Holothurien-Familie Cucumariidae (Ordnung Dendrochirota). 2. Teil: Die Gattungen Cladodaetyla, Hemioedema und Psolidiella. Mitt. hamburg. zool. Mus. 55: 25–38 13 figs.

Patrulius, D. [La position systématique de la forme Pseudosaccocoma strambergensis Remes, Crinoïde aberrant du Malm Supérieur.] Bul. Stiintific. geol. geog. 1 1956: 187–195 5 figs. [In Roumanian with French summary on p. 194.]

Pedrazzi, G. Sull' azione dell' urea nello sviluppo embrionale dei ricci di mare. R.C. Ist. Lombardo, Cl. Sci. Mat. Nat. 91: 672-679 3 figs.

Perlmann, P. (1). Analysis of the surface structures of the sea urchin egg, by means of antibodies. An approach to the study of fertilization. Uppsala (Almqvist and Wiksells 23 pp.

Perlmann, P. (2). Analysis of the surface structures of the sea urchin egg by means of antibodies. I. Comparative study of the effects of various antisera. Exp. Cell Res. 13: 365–390 9 figs.

Perlmann, P. & Hagström, B. E. Cortical inhibitions and polyspermy in antiserum-treated sea urchin eggs. Exp. Cell Res. 12: 418–421.

Perlmann, P. & Perlmann, H. (1). Analysis of the surface structures of the sea urchin egg by means of antibodies. II. The J- and A-antigens. Exp. Cell Res. 13: 454-474 10 figs.

Perlmann, P. & Perlmann, H. (2). Analysis of the surface structures of the sea urchin egg by means of antibodies. III. The C- and F-antigens. Exp. Cell Res. 13: 475–487 7 figs.

Petitot, M.-L. (1). Sur une nouvelle espèce d'echinide irrégulier du Maroc; Pseudopygurus ambroggii nov. sp. Notes Serv. Min. Maroc 121 1954: 83–87 1 pl. 2 figs.

Petitot, M.-L. (2). Sur la présence de Loriolella ludovicii Meneghini dans le Domérian du Moyen Atlas Septentrional. Notes Serv. Min. Maroc 128 1955: 87–94 1 pl.

Petitot, M.-L. (3). Catalogue d'invertébrés fossiles du Maroc. I.Echinides. Notes Serv. Min. Maroc 128 1955: 97-151.

Philip, G. M. Interambulacral Plate Atrophy in *Lovenia woodsi* (Etheridge Fil.). Geol. Mag. 94: 402–408 2 figs.

[Postnikova, I. E., Liashenko, A. I. & Efremova, L. N.] [The stratigraphy of Middle Devonian beds in the Shkapov oil deposits.] C.R. Acad. Sci. URSS 117 2: 275–278 fig. [In Russian.]

Ramsbottom, W. H. C. A rare fossil echinoid from Haw Bank Quarry, Skipton. Naturalist, Lond. No. 862: 106.

Rao, S. R. V. see Gideon, P. W.

Rasmussen, W. H. Cretaceous Crinoidea. Second preliminary report. Medd. dansk. geol. Foren. 12 1954: 553-555.

Raup, D. M. Dendraster: A problem in echinoid taxonomy. J. Paleont. 30 1956: 685-694 4 figs.

Reyment, R. A. Über einige Wirbellose fossilien aus Nigerien und Kamerun, Westafrica. Palaeontographica 109A: 41-70 5 pls. 7 figs.

Reynolds, M. A. The Cainozoic succession of Maslin and Aldinga Bays, South Australia. Trans. roy. Soc. S. Aust. 76 1953: 114–140 3 figs. map.

Ricotta, C. M. B. On the adenosine triphosphatase activity of the particulates of the egg of the sea urchin *Paracentrotus lividus*. Experientia 13:491–492 1 fig.

Roland, J. sec Dieuzeide, R.

Roman, J. Les dyssymétries des *Echinolampas* fossiles. (Échinides exocycles cassiduloïdes.) Bull. Soc. géol. Fr. (6) 7: 67-82 1 fig.

Rothschild, Lord (1). The membrane capacitance of the sea urchin egg. J. biophys. & biochem. Cytol. 3:103-110 l pl.

Rothschild, Lord (2). The fertilising spermatozoon. Discovery 18: 64-65 1 fig.

Rulon, O. Developmental modifications in the sand dollar caused by cobaltous chloride in combination with sodium selenite and zinc chloride. Biol. Bull., Woods Hole, 113: 480-487 33 figs.

Runnström, J. (1). The cell surface in relation to fertilization. Symposia Soc. exp. Biol. No. 6 1952: 39-88 2 pls. 8 figs.

Runnström, J. (2). Cellular structure and behavior under the influence of animal and vegetal factors in sea urchin development. Ark. Zool. 10 523–537 11 figs.

Runnström, J. (3). On the effect of porphyrexid and porphyrindin on the fertilization of the sea urchin egg. Exp. Cell Res. 12: 374–394 15 figs.

Runnström, J. (4). Surface layers of the sea urchin egg and their role in fertilization. Festschrift Arthur Stoll, Basel (Sandoz Ag.): 850-867 11 figs.

Runnström, J., see Kriszat, G.

Runnström, J. & Kriszat, G. The solidifying effect of certain oxidation-reduction indicators on the surface of the unfertilized sea urchin egg. Exp. Cell Res. 12: 526–536 3 figs.

S., S. V. O. Beche-de-mer—the sea's strangest animal. Loris **7** 1955: 82–83 1 fig.

Sagara, J. & Sugawara, K. On the optimum temperature for the development of the starfish, Asterias amurensis. Bull. Tokai. Fish. Res. Lab., No. 16: 49-52 2 figs. 2 pls. [Japanese with English summary.]

Sando, W. J. Beekmantown Group (Lower Ordovician) of Maryland. Mem. geol. Soc. Amer. 68: 1-161 15 pls 20 figs.

Sara, M. Problemi di speciazione negli animali marini. Arch. zool. (ital.) Napoli 42 Suppl.: 273–379 21 figs.

Schlieper, C. Comparative study of Asterias rubens and Mytilus edulis from the North Sea (30 per 1,000 S) and the Western Baltic Sea (15 per 1,000 S). Année biol. (3) 33: 117-127 9 figs.

Schmähl, D. see Druckrey, H.

Schneider, C. O. Sobre la distribucion geografica del *Heliaster helianthus* (Lamk.). Rev. Univ. Santiago **29** (An. Acad. chil. Cienc. nat. No. 9) 1944: 75–76.

Seck, C. Untersuchungen zur Frage der Ionenregulation bei in Brackwasser lebenden Evertebraten. Kieler Meeresforsch. 13: 220–243.

Shaw, A. B. Paleontology of Northwestern Vermont. VI. The Early Middle Cambrian Fauna. J. Paleont. 31: 785–792 2 pls.

Sieverts-Doreck, H. (1). Neufunde von Diamenocrinus und Ctenocrinus aus der Siegen-Stufe des Siegerlandes. Notizbl. hess. Landesamt. Bodenforsch. 85: 62-66 1 fig.

Sieverts-Doreck, H. (2). Bemerkungen über altpaläozoische Crinoiden aus Argentinien. Neues Jb. Geol. Paläont. B. 1957: 151–156 4 figs.

Sinclair, G. W. Date of some species of *Pentremites*. J. Paleont. **31**:982.

Stearn, C. W. A new echinoid from the upper devonian of Alberta. J. Paleont. 30 1956: 741-746 1 pl. 2 figs.

Sternberg, G. F. see Miller, H. W.

Stevenson, R. E. Observations on the morphology of *Astrodapsis*. J. Paleont. **30** 1956: 338-344 1 pl. 1 fig.

Stott, F. C. Observations on the food canal and associated structures in the holothurian *Holothuria forskali* Delle Chiaje. Proc. zool. Soc. Lond. 129: 129–136 5 figs.

Stresemann, E. & others (incl. Gruner, H. E.). Exkursionsfauna von Deutschland. Wirbellose I. Berlin, (Volk & Wissen Volks. Verlag) pp. xxxii + 488, 467 figs. (Echinoderma on pp. 452-460).

Strimple, H. L. Two aberrant crinoid specimens. J. Wash. Acad. Sci. 47: 369 1 fig.

Strimple, H. L. & Koenig, J. W. Mississippian microcrinoids from Oklahoma and New Mexico. J. Paleont. 30 1956: 1225–1247 4 figs.

Struve, W. Ein Massengrab kreidezeitlicher Seelilien. Die *Uintacrinus*-Platte des Senkenberg-Museums. Natur u. Volk 87: 361–373 14 figs.

Sugawara, K. see Sagara, J.

Swann, M. M. The nucleus in fertilization mitosis and cell-division. Symposia Soc. exp. Biol No. 6 1952: 89–104 1 fig.

Sylvester-Bradley, P. C. & Hodson, F. The Fuller's Earth of Whatley, Somerset. Geol. Mag. 94: 312-322.

Szörényi, E. Echinodermenreste aus dem Strážov-Gebirge und aus dem Slowakischen Paradies. Geol. Práce Zprávy Slovensk. Akad. 11: 129–134 1 pl.

Teixeira, C. A propósito da descrição de um novo equinóide fóssil de Angola. Naturalia 6 1956: 83–84.

Termier, G. see Choubert, H.

Thompson, M. S. see Karnovsky, M.

Tommasi, L. R. Os equinodermas do litoral de São Paulo. I. Echinoidea, Crinoidea e Holothuroidea do bentos costeiro. Pap. Dep. Zool. Sec. Agric. S. Paulo 13: 19-44.

Tortonese, E. (1). Intorno ai clipeastroidi del gen. *Rotula*. Doriana 2 (74): 1-4.

Tortonese, E. (2). Variabilità infraspecifica in una popolazione di *Leptasterias polaris* (Müll. Trosch.). Doriana 2 (77): 1–5 2 pls.

Tortonese, E. (3). On the echinoderm fauna of Haifa Bay. Bull. res. Counc. Israel 6B: 189-192.

Tortonese, E. (4). Elementi termofili nell' asterofauna del Mar Ligure. Ann Mus. Storia nat. Genova 69 1957: 94–98.

Tyler, A. & Brookbank, J. W. (1). Antisera that block cell division in developing eggs of sea urchins. Proc. nat. Acad. Sci. Wash. 42 1956: 304–308.

Tyler, A. & Brookbank, J. W. (2). Inhibition of division and development of sea-urchin eggs by antisera against fertilizin. Proc. nat. Acad. Sci. Wash. 42 1956: 308-313 1 fig.

Ubisch, L. von (1). Über Seeigelmerogone II. Pubbl. Staz. zool. Napoli 30: 279–308 29 figs.

Ubisch, L. von (2). Merogone und Bastarde von Seeigeln. Nova Acta Leop. N.S. 19 129: 1-22 19 figs.

Ubisch, L. von (3). Doppelbildungen bei Seeigeln (Psammechinus microtuberculatus, Paracentrotus lividus). Zool. Anz. 159: 1-12 8 figs.

Urbani, E., see Cotronei, G.

Vevers, H. G. & Millott, N. Carotenoid pigments in the integument of the starfish *Marthasterias glacialis* L. Proc. zool. Soc. Lond. 129: 75-80 1 fig.

[Vialov, O. S.] [Palaeogene of the Curendagh type in Turkmen.] Bull. Soc. Nat. Moscow NS 62 Geol. 32 6: 81-92 2 figs. [In Russian.]

Vokes, H. E. (1). Geography and geology of Maryland. Bull. Md. Dep. Geol. 19: xiv, 1-243, 28 pls. figs.

Vokes, H. E. (2). Miocene fossils of Maryland. Bull. Md. Dep. Geol. 20: vii, 1-85 31 pls. (Echinoderma on p. 39).

Walker, M. V. see Miller, H. W.

Whiteley, A. H. see Bolst, A. L.

Wilhelmi, G. Untersuchungen zur Frage einer zytostatischen Wirkung der Antipyretica anhand von Eientwicklungstests. Pubbl. Staz. Zool. Napoli 30: 108–175.

Woo, P.-L. see Chang, F.-Y.

[Yakovlev, N. N.] (1). Echinodermata, Blastoidea in [Librovitch, L. Atlas of the leading forms of the fossil fauna of the USSR. Vol. 4. Lower Carboniferous.] Leningrad 1941 pp. 70-72 pl. X (pt.). [In Russian.]

[Yakovlev, N. N.] (2). Echinodermata in Nalivkin, D. Atlas of the guide forms of the fossil faunas of the USSR. Vol. 3 Devonian Leninghrad 1947 pl. x pp. 55-57.

[Yakovlev, N. N.] (3). [On the existence in the upper silurian and lower devonian of the U.S.S.R. of sea lilies of the family Crotalocrinitidae.] Annu. Soc. paléont. russe 13 1949: 14-23 2 pls.

[Yakovlev, N. N.] (4). [On discoveries of loboliths in the U.S.S.R. and their biological significance.] Annu. Soc. paléont. russe 14 1953: 18–37 3 pls. 4 figs.

[Yakovlev, N. N.] (5). [On a revision of the characters of the genus *Ristna-crinus* Öpik.] Annu. Soc. paléont. russe 15 1956: 155–157 2 figs.

[Yakovlev, N. N.] (6). [Two evolutionary lines of Cromyocrinid sea lilies in connection with their geographical distribution.] [Trans. I. Sess. Russ. Paleont. Soc.] 1955 1957; 11–141 pl.

Yamasu, T. see Kawaguti, S.

Yasumasu, I. see Ishida, J.

Yoshida, M. Positive phototaxis in *Psammechinus microtuberculatus* (Blainville). Pubbl. Staz. zool. Napoli 30: 260–262 1 fig.

Yoshida, M., see Millott, N.

Zavodnik, D. [A tetramerous Anseropoda placenta Pennant from the Adriatic Sea.] Biol. Vestnik, Ljubljana 5 Biol. sekc. 1956: 121–122 1 fig. [In Serbo-Croat.]

Zbyszewski, G. Le Burdigalien de Lisbonne. Comun. Serv. geol. Portug. **38**: 91-226 19 pls.

Zeuthen, E., see Holter, H.

II.—SUBJECT INDEX

GENERAL LITERATURE AND HISTORY

General works.—German littoral echinoderms, Gruner in Stresemann.

Textbooks.—Keys to fossils, Chavan & Cailleux; Guthrie & Anderson.

Bibliography.—Of marine ecology: Crinoids, A. M. Clark in Hedgreth, Holothurians, Asteroids and Holothurians, Deichmann in Hedgreth, Echinoids, Cooke in Hedgreth; of marine paleoecology: Post-Paleozoic Asterozoa, Berry in Ladd, Blastoids, Cline & Beaver in Ladd, Post-Paleozoic Echinoids, Cooke in Ladd, Paleozoic Asterozoa and Echinoids, Cooper in Ladd, Holothurians, Frizzell & Exline in Ladd, Crinoids, Laudon in Ladd, Cystoids, Sinclair in Ladd.

Taxonomy.—Direction 72; Direction 73; of echinoids, Durham & Melville (1) & (2); of Dendraster, Raup; dates of some Pentremites spp., Sinclair; of Rotula, Tortonese (1).

Collections.—Oceanographic Institute of San Pablo University, Bernasconi (1).

STRUCTURE

Symmetry.—Asymmetry in fossil Echinolampas, Roman; double forms of echinoids, Ubisch (3); tetramerous Anseropoda, Zavodnik.

Skeletal anatomy.—Pentremites, Galloway & Kaska; echinoids, Kier (1); interambulacral plate atrophy in fossil Lovenia, Philip; Scyphocrinus with its lobolith root or Camarocrinus, Yakovlev (4).

Internal anatomy.—Cuvierian tubules of *Holothuria*, Endean (1); retraction of starfish stomach, Basch; of Scutellids, Stevenson; food canal in *Holothuria*, Stott.

Microstructure.—Plates of carpoid Paranacystis, Caster & Eaton; Cuvierian tubules of Holothuria, Endean; Vitelline body in Arbacia eggs, Millonig; Cell membranes, Mitchison; microstructure of developing eggs, Runnström (2).

PHYSIOLOGY

Musculature.—Contraction of fibres retracting starfish stomach, Basch.

Reaction to light.—Sensitivity, Millott (2); spectral sensitivity, Millott & Yoshida; positive phototaxis in Psammechinus, Yoshida.

Respiration.—Measurements in early development, Holter & Zeuthen; in Spermatozoa, Mohri (2).

Relation to water.—Quality of seawaters estimated by echinoid development, Bernhard; variations in Asterias according to salinity, Schlieper; ion regulation in brackish water forms, Seck.

Pigments.—Carotenoid in Peronella, Kawaguti & Yamasu; Naphthaquinone pigment in Diadema, Millott (1); Echinochrome A, Nishibori; carotenoids in starfish integument, Vevers & Millott. Bio-chemistry.—Carbonates in relation to skeleton, Revelle and Fairbridge in Hedgpeth, chapter 10; lipids of pyloric cecum of Asterias, Karnovsky, Jeffrey, Thompson & Deane; systematic serology of four asteroids, Kubo; Naphthaquinone pigment in Diadema, Millott (1).

REPRODUCTION AND SEX

Reproductive cycles.—Giese et al. Spermatogenesis.—In Asterina, Cognetti.

Hermaphroditism. — In Echinaster sepositus, Delavault & Cognetti.

DEVELOPMENT

Embryology.—Gametes studied by electron microscopy, Afzelius; dynamics of early development, Holter & Zeuthen; Cleavage mechanism, Mitchison; development of Hemicentrotus, Okada & Miyauchi; mitosis and cell division, Swann.

Fertilization including experimental studies.-Acrosomal reaction, Afzelius (2): partial fertilization induced by oestradiol, Agrell (2); Spermatogenesis and fertilization in Asterina, Cognetti: acrosome reaction, Dan; polyspermy, Hagström: fertilization of jelly-free echinoid eggs, Hagström & Markman; cytochemical studies of fertilization, Immers; aldolase at fertilization, Ishihara: fertilization-promoting factor in egg water, Lybing & Hagström; interaction between sperm and jelly-coat, Monroy (1); swelling properties of egg mitochondria before and after fertilization, Monroy (2); Colleterin in toughening of fertilization membrane, Motomura: analysis of egg surface structures using antibodies, Perlmann (1) & (2), Perlmann & Perlmann (1) & (2); surface folding of unfertilized egg and increased membrane capacitance after fertilization, Rothschild (1); Rothschild (2); cell surface in relation to fertilization, Runnström (1); effects of porphyrexid and porphyrindin on fertilization, Runnström (3); role of surface layers of eggs in fertilization, Runnström (4); solidifying effect on egg surface of porphyrindin and trypsin, Runnström & Kriszat.

Chemical embryology. - Basophilic structures in egg cytoplasm, Afzelius (1): ascorbic acid occurrence in embryos, Bäckström; occurrence of amine oxidase, Blaschko & Hope: phosphorus metabolism in development. Bolst & Whiteley: alkaline phosphatase in hydrid echinoid embryos, Flickinger; acid-soluble nucleotides in early development. Hultin (1): metabolic utilization of C14-formate in embryos, Hultin (2): polysaccharides in mitotic figures. Immers: phospholipid in spermatozoa, Kinoshita, Mohri (2): glucose oxidation traced by radioactive CO2, Krahl; cytochrome oxidase distribution in echinoid larva. Mancuso: thymine in the acid-soluble fraction of Arbacia eggs. Marshak & Marshak: perivisceral fluid of Arbacia eggs, Messina; fatty acid oxidation in spermatozoa, Mohri (1): effect of malonate on O2 uptake and dehydrogenase activity of spermatozoa, Mohri (3); ATP-ases of mitochondria, Ricotta.

Experimental embryology.—Effects of temperature changes on cellular division, Agrell (1); mass separation of developing blastula fragments, Banhidi; 'vitellogenous band' of oocyte, Cotronei & Urbani; inhibitors of cell division in unfertilized echinoid eggs. Druckrey, Schmähl & Kaiser; effect of toluidine blue on starfish egg. Fautrez & Fautrez-Firlefyn; effects of amino acids, Gustafson & Hörstadius; determination of bilateral symmetry in eggs, Hörstadius (1): regulation of bilateral symmetry in split and fused 32-cell eggs, Hörstadius (2); uptake of Ca45 by echinoid eggs, Hsaio & Boroughs; changes in protein specificity, Ishida & Yasumasu; effect on cleavage of demecolcin, Kobayashi & Nakamura (1) & (2): effect of DNP, Kriszat & Runnström; effect of different physiological conditions on nuclear size, Kupka; animalization by certain polysulphonic derivatives, Lallier (1); effect of perchlorate on determination, Lallier (2); animalization by sodium polyanethol sulphonate, Lallier (3); effects of three dyes on determination, Lallier (4), of sulphydrilic substances, Lallier (5), and of sulphonic derivatives on determination, Lallier (6); nuclear activity in differentiation, Markman; protoplasmic contractility and

structure, Marsland; morphogenetic effects of treatment of eggs with trypsin, Moore (2); Infraciliature in Paracentrotus | arvae, Mugard; incorporation of radioactive isotopes into echinoid eggs, Nakano & Monroy; effect of urea on development, Pedrazzi; analysis of egg surface structures using antibodies, Perlmann (1) & (2) and Perlmann & Perlmann (1) & (2); cortical inhibitions and polyspermy in antiserum-treated eggs, Perlmann & Hagström; effects of cobaltous chloride with sodium chloride and zinc chloride on development of sand-dollar larva, Rulon; effect on structure of animal and vegetal factors in echinoid development, Runnström (2); antisera blocking cell division, Tyler & Brookbank (1) & (2); abnormal and hybrid larvae, Ubisch (1) & (2); cytostatic effect of antipyretics on egg development, Wilhelmi.

Techniques.—Electron microscopy in studying gametes, Afzelius (3) & (4) and for Arbacia eggs, Millonig; phase contrast microscopy for observing development, Holter & Zeuthen.

EVOLUTION AND GENETICS

Phylogeny.—Of groups of Pentremites, Galloway & Kaska.

Speciation.-Sarà.

Hybridization. — Biparental inheritance in interordinal echinoid crossing, Moore (1); Hybrid larvae, Ubisch (2).

Variation.—In Echinocyamus, Kier (4), in Dendraster, Raup, in Leptasterias polaris, Tortonese (2).

ECOLOGY AND HABITS

General works.-Hedgpeth et al.

Paleoecology.—Martinsburg shale, Pennsylvania, U.S., Cramer; of Pentremites, Galloway & Kaska; general, Ladd et al.; of Saccocoma, Patrulius.

Occurrence.-Beche de mer, S.V.O.S.

Communities.—Of fishery grounds in Channel and North Sea, Ancellin; fluctuations in populations of Asterias, Coe; Zonation, Collignon; bottom communities, Thorson in Hedgpeth; starfishes in Ligurian Sea, Tortonese (4).

Effects of environment.—Distribution of larvae around Kiel, Banse; optimum temperature for *Asterias* development, Sagara & Sugawara; salinity differences and *Asterias*, Schlieper.

Feeding.—Of Evasterias, Christensen, of Psammechinus in aquaria, Hancock.

Parasites.—Umagillidae (Turbellarians) forming 'brown bodies' in *Holothuria*, Arvy.

Commensals.—Some known examples, Dales in Hedgpeth.

Aquaria.—Psammechinus, Hancock.

DISTRIBUTION

GEOGRAPHICAL

Arctic.—Bering Sea, Baranova; Spitzbergen, Blacker; Alaska, Durham.

North Atlantic.—English Channel and North Sea, Ancellin; Pembrokeshire, Wales, Bassindale & Barrett; North Carolina, U.S., Cooke (2); Portugal, Astropecten, Cumano; Maine, U.S., Frings & Frings; Germany, Gruner in Stresemann; SW. England, Marine Biological Association.

Mediterranean.—Cherbonnier (1); Algeria, Dieuzeide & Roland; Gautier-Michaz; Israel, Tortonese (3); Ligurian Sea, Asteroids, Tortonese (4).

Tropical Atlantic.—Sierra Leone, Cherbonnier, (3); Congo, Collignon; Bahama holothurians, Deichmann; Tropical West Africa, Madsen (3).

Indo-West Pacific.—Portuguese E. Africa, Balinsky; N. China, Chang & Woo; Djakonov; Guam, Holothurians, Domantay; Queensland, Australia, Endean (2); Gulf of Cutch, India, Gideon, Menon, Rao & Jose; Japan, Asteroids, Hayashi (1) & (2).

East Pacific.—Clipperton Island, Hertlein & Emerson; S. America, especially Chile, Madsen (1); Heliaster, Schneider.

South Temperate and Subantarctic.— South-east Africa, Balinsky; east S. America, Bernasconi (1); Brazil, Asteroid, Bernasconi (2); Patagonia, Madsen (1); Brazil, Tommasi.

GEOLOGICAL

General.—Marvland, U.S., Vokes (1),

Pleistocene.—Mexico, Clypeaster, Caso (1) & (2); Maastricht, Engel in Burck; Holothurians, Frizzell & Exline; Gulf of California, Hertlein; Mariana Is., Johnson; British Somaliland, Kier (4); S. Australia, Reynolds.

Pliocene.—Mexico, Clypeaster, Caso (1) & (2); Alaska, Durham; Maastricht, Engel in Burck; holothurians, Frizzell & Exline; Gulf of California, Hertlein; Morocco, Echinoids, Petitot (3); S. Australia, Reynolds.

Miocene.—Mexico, Clypeaster, Caso (1) & (2); Mariana Is., Cooke (1), Johnson; Central America and U.S., Durham; holothurians, Frizzell & Exline; Germany, Jörg; British Somaliland, Kier (4); Italy, Mirigliano; Japan, Morishita (1) & (2); Morocco, Echinoids, Petitot (3); Astrodapsis, Stevenson; Maryland, U.S., Vokes (2); Portugal, Zbyszewski.

Oligocene.—Mexico, Clypeaster, Caso (1) & (2); Oregon, U.S., Durham; holothurians, Frizzell & Exline; Italy, Mirigliano; Japan, Morishita (1); Rasmussen; S. Australia, Reynolds; Turkmen, U.S.S.R., Vialov.

Eocene.—Spain, Bataller; holothurians, Frizzell & Exline; Mariana Is., Johnson; British Somaliland, Kier (4); Morocco, Echinoids, Petitot (3); S. Australia, Reynolds; France, Roman; Turkmen, U.S.S.R., Vialov.

Cretaceous.—Azerbaijan, U.S.S.R., Aliev; Spain, Bataller; E. Greenland, Donovan; Maastricht, Engel & Meijer; holothurians, Frizzell & Exline; Egypt, Jux; Uintacrinus, Miller, Sternberg & Walker; Japan, Morishita (1); Morocco, Echinoids, Petitot (3); Crinoids, Rasmussen; Nigeria and Cameroons, Reyment; Germany, Struve; Slovakia, Szörényi.

Jurassic.—Algeria, Devriès; E. Greenland, Donovan; holothurians, Frizzell & Exline; Dorset, England, holothurians, Hampton; Dorset, Hodson, Harris & Lawson; Pomerania, Kongiel; Roumania, Patrulius; Morocco, Petitot (1) & (3); Algeria, Petitot (2); Somerset, England, Sylvester-Bradley & Hodson.

Triassic.—Gərmuny, Həllər; Morocco, Echinoids, Petitot (3); Slovakia, Szörén-yi.

Permian.—Cromyocrinidae, Yakov-lev (6).

Carboniferous. — Kentucky, U.S., Conkin; Nevada, U.S., Durham; U.S.S.R. echinoids, Faas; N. America, Pentremites, Galloway & Kaska; Ireland, echinoid, Kier (2); Texas, U.S., Kier (3); Arizona, U.S., holothurians, Langenheim & Epis; Morocco, echinoids, Petitot (3); Yorkshire, Ramsbottom; Oklahoma, U.S., Strimple; Oklahoma and New Mexico, U.S., Strimple & Koenig; U.S.S.R. blastoids, Yakovlev (1); Cromyocrinidae, Yakovlev (6).

Devonian.—Brazil, Caster & Eaton; U.S.S.R., Eltuishcheva; Rhineland, Germany, Falke; E. France, Firtion; Victoria, Australia, Carpoids, Gill & Caster; Germany, Grabert & Grabert; Germany, Asterozoa, Lehmann; Bashkir, U.S.S.R., Crinoids, Postnikova, Liashenko & Efremova; Germany, Sieverts-Doreck (1); Alberta, Canada, Stearn; U.S.S.R., Yakovlev (2) & (3).

Silurian. — U.S.S.R., Eltuishcheva; Victoria, Australia, Carpoids, Gill & Caster; U.S.S.R., Yakovlev (3) & (4).

Ordovician. — Oklahoma, Amsden; Morocco, Choubert & Termier; Pennsylvania, U.S., Asterozoa, Cramer; U.S.S.R., Eltuishcheva, Ghekker & Ghekker; Maryland, U.S., Sando; Argentina, Sieverts-Doreck (2); Estonia, Yakovlev (5).

Cambrian.—Walcott's 'Holothurians,' Madsen (2); Vermont, U.S., Shaw; Argentina, Sieverts-Doreck (2).

III.—SYSTEMATIC INDEX

HOLOTHURIOIDEA

†Achistrum cordatum sp. nov. Jurassio Dorset, England, Hampton p. 509 fig. 9, A. cf. bathonianum figs. 3, 4, A. gamma fig. 5, monochordata fig. 6, A. sp. ind. figs. 12, 13 monochordata pp. 340-341 figs. 10, 11, spp. nov. Jurassic Dorset, England, Hodson, Harris & Lawson, A.

cf. issleri p. 340 figs. 8, 9, Id.; A. frizzelli sp. nov. Carboniferous Arizona, U.S., Langenheim & Epis pp. 167–170 pl. i figs. 1-6.

Actinopyga agassizii, Deichmann pp. 6-7 figs. 5-13; A. mauritiana, Domantay pp. 349-350.

Capheira mollis, Baranova p. 239.

Chiridota discolor, pellucida, p. 245, Baranova.

†Chiridotitesidae, nom. nov. for family Theeliidae Frizzell & Exline, invalidated, Deflandre-Rigaud p. 354.

Cladodactyla pp. 25–27, with C. crocea pp. 27–29 figs. 10–13, senegalensis pp. 29–31 figs. 14–19, Panning.

†Croneisites gen. nov. pp. 113-114, family Synaptitidae, for type Synapta oligocaenica Spandel, Frizzell & Exline; C. laevigatus, oligocaenicus p. 114, Id.

Cucumaria abyssorum pp. 241–242, calcigera p. 241, californica p. 242, chronchjelmi p. 240, falax p. 242, glacialis p. 240, ijimai p. 242, japonica p. 240, lamperti p. 242, pusilla p. 241, vegae p. 241, Baranova; C. chronhjelmi, Chang & Woo (in Chinese) pp. 137–139 fig. 9 pl. ii figs. 3–4; C. adela p. 36, frauenfeldi pp. 36–37, posthuma p. 36, sancti-johannis p. 37, Panning.

†Eldonia ludwigi, Madsen (2) p. 281, a coelenterate.

Elpidia glacialis, Baranova p. 236. †Etheridgella sp., Hampton p. 510 figs. 11, 12.

Euapta lappa, Deichmann p. 17; E. godeffroyi, Domantay p. 353.

Hemioedema pp. 32–34, with H. spectabilis pp. 34–35 figs. 20–22, Panning.

Holothuria Linnaeus 1758, suppressed, Direction 72 p. 169; H. tremula Linnaeus 1767 placed on the Official List of Specific Names in Zoology, Direction 73 pp. 198, 208; H. floridana figs. 39-53, glaberrima p. 12, grisea pp. 11-12, impatiens pp. 7-8 figs. 14, 15, mexicana pp. 9-11 figs. 21-38, parvula pp. 12-13, princeps p. 8 figs. 16-20, Deichmann; H. arenicola pp. 346-347, argus pp. 338-339, atra pp. 339-340, cinerascens pp. 340-341, curiosa var. pervicax pp. 341-342, edulis pp. 342-343, gyrifer pp. 343-344, impatiens pp. 344-346, pulla pp. 347-348, rugosa p. 348, sanctori

pp. 348-349, Domantay; H. leucospilota, Hertlein & Emerson pp. 6-7; H. grisea, Tommasi pp. 40-41 figs. 28, 30 d, e pl. iv fig. 1.

†Laggania cambria, Madsen (2) p. 281 a polychaete.

Leptosynapta ooplax, Chang & Woo pp. 141-143 fig. 11; L. inhaerens, Gruner in Stresemann p. 460 fig. 460/1.

†Louisella pedunculata, Madsen (2) p. 281, a polychaete.

†Mackenzia costalis, Madsen (2) p. 281, a coelenterate.

Myriotrochus rinkii, **Baranova** pp. 245–246.

Opheodesoma spectabilis, Domantay p. 354.

Paelopatides solea, Baranova pp. 238–239 fig. 19.

Pannychia moseleyi virgulifera, Baranova pp. 235–236.

Parathyone gen. nov., family Cucumariidae, for type Thyone surinamensis Semper, Deichmann p. 16; P. surinamensis p. 16.

Pentacucumis gen. nov., family Cucumariidae, for type Cladodactyla planci Brandt, Deichmann p. 13; P. planci pp. 13-16 figs. 54-71, Id.

Phyllophorus glaucus, Baranova p. 243; P.ordinatus, Chang & Woo pp. 139–140 fig. 10 pl. ii fig. 5.

†Portalia mira, Madsen (2) p. 281, a sponge.

Pseudostichopus aleutianus p. 239, nudus p. 240, trachus p. 239, Baranova.

Psolidiella, Panning p. 35.

Psolidum bullatum, Baranova p. 245.

Psolus chitonoides pp. 244-245, fabricii p. 244, peronii p. 244, regalis p. 243, Baranova

† Redoubtia polypodia, Madsen (2) p. 281, a polychaete.

†Rhabdotites bifidus p. 342 figs. 16, 22, divergens p. 342 figs. 14, 17, 18, dorsetensis p. 341 figs. 5, 15, 20, 21, irregularis p. 343 fig. 25, tridens p. 343 fig. 19, spp. nov. Jurassic Dorset, England, Hodson, Harris & Lawson; R.? rectus, Hampton p. 510 fig. 10.

†Rigaudites gen. nov. pp. 101-102, family Synaptitidae, for type Synaptites cuvillieri Deflandre-Rigaud, Frizzell & Exline; R. bastropanus pp. 103-104 fig. 5 Eccene Texas, U.S., plummerae pp. 106-107 fig. 3 Cretaceous Texas, spinosus pp. 107-108 fig. 4 Cretaceous Texas, tallaili pp. 108-110 fig. 6 Cretaceous Texas spp. nov., R. sp. nov. Cretaceous Texas pp. 110-111, cuvillieri pp. 105-106 figs. 1 2, Id.

†Rota gen. nov. p. 170, family Theelidae, for type R. martini sp. nov. Carboniferous Arizona, U.S. p. 170 pl. i figs. 17–25, Langenheim & Epis.

Scotoplanes murrayi, Baranova p. 237.

†Spandelites gen. nov. p. 101, family Synaptitidae, for type Synaptites? irregularis Deflandre-Rigaud, Frizzell & Exline.

Sphaerothuria bitentaculata, Baranova pp. 242–243.

Stichopus japonicus v. armatus, Baranova p. 237; S. badionotus, Deichmann pp. 4-5 figs. 1-4; S. badionatus pp. 350-351, chloronotus pp. 351-352, horrens p. 352, Domantay; S. badionotus, Tommasi pp. 41-42 fig. 30a pl. iv figs. 3, 4.

Synallactes nozawai, Baranova p. 237.

Synapta maculata, Domantay pp. 353-354.

†Synaptites emended pp. 110-111, with S. circularis pp. 111-112, eocoenus, renifer p. 112, stueri pp. 112-113 truncatus p. 113, Frizzell & Exline.

Synaptula hydriformis, Tommasi pp. 39–40 fig. 30 b, c pl. iv fig. 2.

†Theelia Schlumberger invalidated, and a synonym of Chiridotites Deflandre-Rigaud, Deflandre-Rigaud, pp. 353-354; T. wessexensis sp. nov Jurassic Dorset, England, Hodson, Harris & Lawson pp. 338-339 figs. 1-3, 5-7, 17 T. convexa fig. 4, Id.

†Theeliidae Frizzell & Exline invalidated, replaced by Chiridotitesidae, Deflandre-Rigaud.

†Thuroholia with T. marginata sp. nov. Carboniferous Arizona, U.S., Langenheim & Epis p. 167 pl. i figs. 7-16.

Thyonacta sabanillaensis, Cherbonnier (2) pp. 537-540 figs. 1, 2.

Thyone fusus, Gruner in Stresemann p. 460 fig. 459 3; T. briareus, Guthrie & Anderson p. 503 figs. 16.12, 16.13.

Thyonidium pellucidum, Baranova p. 243.

ECHINOIDEA

†Abertella kewi Miocene Mexico pp. 627-628 fig. 2 pl. lxxii figs. 1, 7, palmeri? Miocene Guatemala pp. 626-627 pl. lxxii figs. 4, 9. 10, spp. nov., Durham; A. aberti, Vokes (2) p. 39 pl. xxx figs. 4, 5.

†Acanthocidaris sp., Cooke (1) p. 362 pl. exix fig. 8.

†Acrosalenia angularis, Kongiel p. 60 pl. i figs. 1-3.

Aëropsis fulva, Baranova pp. 232–234 fig. 18.

†Albertechinus gen. nov. pp. 741-744, family Lepidocentridae, for type A. montanus sp. nov. Devonian Alberta, Canada pp. 744-746 fig. 1 pl. lxxxi figs. 1-3, Stearn.

†Ambipleurus rotundatus sp. nov. Eocene British Somaliland Kier (4) p. 847 fig. 1e-g pl. ciii fig. 9; A. douvillei pp. 847-848 fig. 1c, d pl. ciii fig. 10, Id.

†Amphiope bioculata, Chavan & Cailleux p. 199 fig. 2245.

Aporocidaris fragilis, Baranova pp. 209–210.

Arbacia lixula, Bernasconi (1) pp-55-56 pl. i figs. 3, 4 pl. iv fig. 3; Alixula var. africana, Collignon pl. ii fig. 3B; A. punctulata, Guthrie & Anderson fig. 16.9 A, B; A. lixula, Tommasi pp. 23-25 figs. 8-11 pl. i fig. 1.

Arbacioida, Durham & Melville (2) p. 255, diagnosed.

†Arcaechinus gen. nov. p. 891, family Brissidae for type A. auraduensis sp. nov. Eccene British Somaliland pp. 891-893 fig. 18 pl. cvii figs. 6-8, Kier (4).

 $\dagger Archaeocidaris$ sp., Faas p. 75 pl. x fig. 10.

†Astrodapsis pp. 340-343 with A. cuyamanus fig. 1E, peltoides pl. xlv fig. 6, woodringi pl. xlv fig. 7, Stevenson.

Atelostomata, Durham & Melville (2) p. 259, diagnosed.

†Bonaireaster rutteni, Durham & Melville (2) figs. 3, 4.

†Bothriocidaris, Durham & Melville (2) p. 242.

†Bothriocidaroida, **Durham & Mel-ville** (2) p. 249, diagnosed.

†Brightonia gen. nov. pp. 871–872, family Stenonasteridae, for type B. macfadyeni sp. nov. Paleocene British Somaliland pp. 872–873 fig. 13 pl. civ figs. 12, 13, Kier (4).

Brisaster latifrons p. 235, townsendi pp. 234-235, Baranova.

Brissopsis lyrifera, Gruner in Stresemann p. 459 fig. 459/2, Madsen (3) p. 479.

†Brissopsis makiyamai sp. nov. Miocene Japan, Morishita (2) pp. 161-163 pl. i figs. 1-5; B. sp. cf. raulini, Kier (4) p. 899 pl. evii figs. 16, 17.

Cassiduloida, **Durham & Melville** (2) p. 260, diagnosed.

Centrostephanus besnardi, Bernasconi (1) pp. 56-59 text figs. 1-9 pl. i figs. 2, 6, pl. vii fig. 2a-c; C. longispinus, Millott (2) fig. 4.

Cidaris cidaris nuda, Madsen (3) pp. 479-481.

†Cidaris coronata, Chavan & Cailleux p. 193 fig. 2220; C. dorsata, Szörényi pp. 130–131.

Cidaroida, **Durham & Melville** (2) p. 251, diagnosed.

Clypeaster speciosus, Stevenson fig. 1 A, B pl. xlv figs. 4, 5.

Clypeaster (Stolonoclypus) subdepressus, Tommasi pp. 30–31 figs. 22–24 pl. ii figs. 3, 4.

†Clypeaster saipanicus sp. nov. Miocene Mariana Is., Pacific, Cooke (1) p. 362 pl. cxix figs. 14-17; C. chiapasensis pp. 502-505 figs. 4, 5, C. aff. chiapasensis pp. 519-521 fig. 17, C. crustulum pp. 513-516 figs. 12, 13, marinanus pp. 508-509, meridanensis pp. 511-513 fig. 11, oxybaphon pp. 505-508 figs. 7, 8, pileus pp. 516-519 figs. 14-16, speciosus pp. 521-524 fig. 18, staubi pp. 499-502 figs. 1-3, topilanus pp. 509-511 figs. 9, 10, Caso (2); C. aegypticus, Chavan & Cailleux p. 200 fig. 2249.

Clypeasterina, **Durham & Melville** (2) p. 259, diagnosed.

Clypeasteroida, **Durham & Melville** (2) p. 259 diagnosed.

Vol. 94.

†Clypeus ploti, Chavan & Cailleux p. 202 fig. 2256.

†Codiopsis doma, Chavan & Cailleux p. 197 fig. 2235.

†Collyrites ellipticus, Chavan & Cailleux p. 203 fig. 2263.

†Collyropsis moussoni, Szörényi pp. 131-133.

Conoclypina, **Durham & Melville** (2) p. 256, diagnosed.

†Conoclypus aequidilatatus, Durham & Melville (2) fig. 5; C. p. 858 with C. delanouei pp. 858-859 Kier (4).

Dendraster (recent and fossil) with D. excentricus, Raup pp. 685-694; D. excentricus, Stevenson fig. 1 C, D pl. xlv figs, 2, 3.

Diadema antillarum, Millott (2) figs. 1-3, 13, 15.

Diadematacea, **Durham & Melville** (2) pp. 252-253, diagnosed.

Diadematoida, Durham & Melville (2) p. 253, diagnosed.

†Discoidea karakaschi, Aliev p. 1171 fig. 3; D. sp.? rotula, Szörényi p. 131.

†Discoides cylindricus, Chavan & Cailleux p. 197 fig. 2236.

†Donbassechinus gen. nov. p. 73, family Palaeechinidae, for type D. kumpani sp. nov. Carboniferous Donbasin, U.S.S.R., pp. 73-74 figs. 6-8 pl. x figs. 6-8, Faas.

Echinacea, **Durham & Melville** (2) p. 253, diagnosed.

†Echinanthus vidali, Bataller p. 30 pl. v figs. 3-6.

Echinarachnius asiaticus pp. 228–229 fig. 16, parma pp. 225–228 fig. 15, Baranova; E. parma, Durham pp. 629–630 pl. lxxii fig. 5.

†*Echinarachnius alaskensis* **sp. nov.** Pliocene Alaska, **Durham** pp. 628-629 pl. lxxii figs. 6, 8.

Echinocardium cordatum, Chang & Woo pp. 135-137 fig. 8 pl. i figs. 3-5, Gruner in Stresemann p. 459 fig. 459/1, Kier (1) p. 971 fig. 1.

†Echinocardium orthonotum, Vokes (2) p. 39 pl. xxx figs. 1–3.

†Echinocorys ovatus, Chavan & Cailleux p. 204 fig. 2264.

†Echinocyamus p. 859 with E. polymorpha pp. 859-870 figs. 7-12 pl. civ figs. 5-8, Kier (4).

†Echinocystitoida, Durham & Melville (2) p. 249, diagnosed.

†Echinodiscus chikuzenensis, Morishita (1) p. 194.

Echinoida, Durham & Melville (2) p. 255, diagnosed.

†Echinolampas vidali, Bataller p. 31 pl. v figs. 7-9; E. calvimontanum, Chavan & Cailleux p. 203 fig. 2262; E. sp. Cooke (1) p. 364 pl. cxix figs. 22, 23; E. p. 848 with E. fraasi pp. 848-855 figs. 2-5 pl. ciii figs. 11, 12 pl. civ fig. 1, Kier (4); E. stelliferus, Roman fig. 1.

Echinometra lucunter, Bernasconi (1) pp. 62-63 pl. ii figs. 1, 5, Tommasi pp. 29-30 figs. 16, 20 pl. i figs. 4, 6.

Echinoneina, Durham & Melville (2) p. 256, diagnosed.

†Echinostrephus saipanicum sp. nov. Miocene Mariana Is., Cooke (1) p. 362 pl. exix figs. 4-6.

Echinothurioida, **Durham & Melville** (2) p. 253, diagnosed.

Echinus acutus mediterraneus, Madsen (3) p. 481.

Encope emarginata, Bernasconi (1) pp. 64-65 pl. iii figs. 1, 2, Tommasi p. 34 pl. iii figs. 1, 2; E. californica, Stevenson fig. 1F pl. xlv fig. 1.

†Epiaster ? sp., Reyment pp. 45-46 pl. vii fig. 3a, b.

Eucidaris tribuloides, Bernasconi (1) pp. 52-54 pl. i figs. I, 5, Collignon pl. ii figs. 3A, 4.

Euechinoidea, Durham & Melville (2) p. 252, diagnosed.

†Eupatagus nipponicus sp. nov. Miocene Japan, Morishita (2) pp. 163-164 pl. i figs. 7, 8; E. ornatus, Chavan & Cailleux p. 206 fig. 2274; Eupatagus p. 896 with E. cairensis pp. 896-897 pl. cvii fig. 9, dainellii pp. 897-898 pl. cvii figs. 10-13, fecundus p. 898 pl. cvii figs. 14, 15, E. sp. cf. cordiformis p. 898, Kier (4).

†Eupholidocidaris gen. nov. p. 15 family Lepidocentridae, for type E. brightoni sp. nov. Carboniferous Ireland, pp. 16–17 figs. 1, 2 pl. i, Kier (2); E. belli sp. nov. Carboniferous Texas, Kier (3) pp. 326–328 fig. 1.

†Fibularia dubarensis sp. nov. Miocene British Somaliland Kier (4) pp. 870-871 fig. 7c-e pl. civ figs. 9-11.

†Glypticus hieroglyphicus, Chavan & Cailleux p. 196 fig. 2233.

Glyptocidaris crenularis, Chang & Woo pp. 128-130 figs. 1-3 pl. i fig. 1 pl. ii fig. 1.

Gnathostomata, **Durham & Melville** (2) p. 255, diagnosed.

†Haimea ovumserpentis, Durham & Melville (2) fig. 2.

†Hemiaster bufo, Chavan & Cailleux p. 205 fig. 2269.

†Hemiaster (Trachyaster) with H. (Trachyaster) sp. Kier (4) p. 880.

†Hemicidaris crenularis, Chavan & Cailleux p. 193 fig. 2222.

†Hemicidaris (Hemicidaris) crenularis var. alta var. nov. Jurassic Pomerania, Kongiel pp. 62-63 pl. ii figs. 5-6; H. crenularis pp. 61-62 pl. ii figs. 2-4.

Hemicidaroida, Durham & Melville (2) p. 254, diagnosed.

†Heteraster oblongus, Chavan & Cailleux p. 205 fig. 2270.

†Heterocentrotus sp., Cooke (1) p. 362 pl. cxix fig. 7; H. mammillatus, Kier (4) p. 848.

†Holaster subglobosus, Chavan & Cailleux p. 204 fig. 2266; H. laevis, Szörényi p. 133.

Holasteroida order nov. for families Collyritidae, Disasteridae, Holasteridae, Urechinidae, Calymnidae, Pourtalesiidae and Stenonasteridae, **Durham & Melville** (2) pp. 260–261.

Holectypina, Durham & Melville (2) p. 256, diagnosed.

Holectypoida, Durham & Melville (2) p. 256, diagnosed.

†Holectypus depressus, Chavan & Cailleux p. 197 fig. 2237, Durham & Melville (2) fig. 1A, B; H. corallinus, Kongiel p. 72 pl. vi figs. 3-6.

†Hypodiadema hoffmanni, Kongiel p. 63 pl. iii figs. 1-3.

†Infraclypeus thalebensis, **Devriès** pp 149-157 figs. 1-9.

†Kewia sp. indet., Oligocene Oregon, U.S., Durham p. 630 pl. lxxii fig. 3.

Laganina, Durham & Melville (2) p. 259, diagnosed.

†Leiocidaris thiébaudi, Teixeira p. 83.

†Leviechinus gen. nov. pp. 873-874, family Stenonasteridae, for type Pericosmus gregoryi Currie, Kier (4); L. gregoryi pp. 874-875 fig. 14 pl. civ figs. 14-16, Id.

†Linthia somaliensis, Kier (1) pp. 971–974 fig. 3; Linthia p. 886 with L. cavernosa pp. 888–890 fig. 17 pl. cvi figs. 17–19, somaliensis pp. 886–888 fig. 16 pl. cvi figs. 15, 16, Kier (4); L. praenipponica, Morishita (1) pp. 196–197 pl. ii figs. 2a-d; L. sudanensis, Reyment pp. 47–48 pl. vii figs. 4, 5a, 5b, 6a, 6b.

Loriolella with L. ludovicii, Petitot (2) pp. 87-91 pl. i.

†Lovenechinus sp. cf. septies, Durham pp. 625-626.

†Lovenia woodsi, Philip pp. 402-407 figs. 1, 2.

†Lutetiaster p. 890 with L. maccagnoi pp. 890-891, Kier (4).

†Lychnidius scrobiculatus, Engel & Meijer pp. 88-93 figs. 1-3 pl.

Lytechinus variegatus, Bernasconi (1) pp. 60-61 pl. ii figs. 2, 6, Millott (2) figs. 5, 6, 16-21, Tommasi pp. 25-26 figs. 12-15 pl. i figs. 2, 5.

Megalopoda, **Durham & Melville** (2) p. 262, diagnosed.

Mellita quinquiesperforata latiambulacra, Bernasconi (1) pp. 63-64 pl. iv figs. 5, 6; M. longefissa, Stevenson fig. 1G; M. quinquiesperforata, Tommasi pp. 31-32 fig. 3 pl. ii figs. 1, 2 pl. iii figs. 3, 4.

†Melonechinus (?) heckeri sp. nov. Carboniferous Leningrad Province, U.S.S.R., Faas pp. 74-75 fig. 9 pl. x fig. 9.

Meoma cadenati sp. nov. Senegal, Madsen (3) pp. 474-479 figs. 1-3.

†Merriamaster sp., Stevenson fig. 1H.

Metalia costae, Gauthier-Michaz p. 82.

†Micraster coranguinum, Chavan & Cailleux p. 206 fig. 2271, Kier (1) p. 971 fig. 2.

†Micropedina olisiponensis, Reyment pp. 46-47 fig. la-d pl. vii figs. la-c.

†*Migliorinia* p. 893 with *M. migiurtina* pp. 893–896 figs. 19, 20 pl. cvii figs. 1–5, **Kier** (4).

Miocidaridae, nom. nov. for Streptocidarinae Lambert, inacceptable, Durham & Melville (2) p. 252.

Moira atropos, Bernasconi (1) pp. 65-67 pl. iv figs. 1, 2.

†Myriastiches, Durham & Melville (2) p. 243.

 $Neolampas\ rostellata,\ {f Gautier-Michaz}$ p. 81.

†Nucleolites scutatus, Chavan & Cailleux p. 202 fig. 2257.

Nucleolitoida, **Durham & Melville** (2) p. 260, diagnosed.

†Nudobrissus sp. indet., Morishita (1) p. 200 pl. iv fig. 2.

†Opissaster pp. 875–876 with O. auraduensis sp. nov. Paleocene British Somaliland pp. 876–877 fig. 15a pl. cv figs. 3–5, O. derasmoi var. angulatus var. nov. Eocene British Somaliland pp. 879–880 pl. cv figs. 7–9, Kier (4); O. derasmoi pp. 878–879 fig. 15b pl. cv fig. 6 farquharsoni p. 876, somaliensis pp. 877–878 Id.

†Orthopsis pomeraniae sp. nov. Jurassic Pomerania, Kongiel pp. 70-72 pl. v figs. 5, 6; O. granularis, Durham & Melville (2) fig. 1C.

†Palaechinoida, Durham & Melvill (2) p. 251, diagnosed.

† Palechinus, Chavan & Cailleux pp. 192–193 fig. 2219.

†Palmeraster japonicus sp. nov. Oligocene Japan, Morishita (1) pp. 194–195 pl. ii figs. 1a-c.

Paracentrotus gaimardi, Tommasi pp. 26–29 fig. 19 pl. i fig. 3.

†Parasalenia marianae sp. nov. Miocene Mariana Is., Pacific, Cooke (1) pp. 361–362 pl. exix figs. 1–3.

†Paraster saipanicus sp. nov. Miocene Mariana Is., Pacific, Cooke (1) pp. 363-364 pl. cxix figs. 18-21.

†*Pedinopsis ? melo* sp. nov. Eccene British Somaliland, **Kier** (4) p. 845 fig. 1a, b pl. eiii figs. 4, 5.

†Pericosmus cf. spatangoides, Morishita (1) pp. 195–196 pl. i figs. 2–3.

†Perischocidaris hartiana, Ramsbottom p. 106 fig.

Perischoechinoidea, Durham & Melville (2) p. 249, diagnosed

Peronella japonica, Kawaguti & Yamasu figs. 1-3.

†Pharaonaster with P. sp. cf. ammon Kier (4) p. 875 pl. civ fig. 17 pl. cv figs. 1, 2.

†Phymosoma koenigi, Chavan & Cailleux p. 195 fig. 2227a-c, P. supracorallinum, Kongiel pp. 67-68 pl. iv figs. 4-6.

Phymosomatoida, Durham & Melville (2) p. 254, diagnosed.

Plagiobrissus costae, Madsen (3) p. 479, Tortonese (3) p. 192.

†Plesiaster peini, Morishita (1) pp. 199-200 pl. i fig. 1 pl. iv fig. 3.

†Plesiolampas p. 855 with P. auraduensis pp. 857-858 pl. civ figs. 2-4, curriae pp. 856-857 fig. 6 pl. ciii figs. 13, 14, spp. nov. Paleocene British Somaliland, Kier (4); P. saharae, Reyment p. 47 fig. 1e pl. vii figs. 2a-b.

†Porocidaris schmidelii, Kier (4) p. 844 pl. ciii fig. 3.

†Porosoma vidali, Bataller p. 31 pl. v¹ figs. 1, 1a, b; P. lamberti, Kier (4) p. 845 pl. ciii figs. 6-8.

Pourtalesia laguncula beringiana, Baranova pp. 231–232 fig. 17.

†Pseudocidaris (Pseudocidaris) mammosa, Kongiel pp. 60-61 pl. i figs. 4-6 pl. ii fig. 1.

†Pseudopygurus ambroggii sp. nov. Jurassic Morocco, Petitot (1) pp. 83–86 figs. 1, 2 pl. i figs. 1–7.

†Pseudopyrina ovulum, Chavan & Cailleux p. 201 fig. 2252.

†Pygaster truncatus, Chavan & Cailleux p. 198 fig. 2238.

†Pygaster (Pygaster) cf. subtilis, Kongiel p. 72 pl. vi figs. 1, 2.

Pygasteroida order nov. for family Pygasteridae Lambert, 1900, **Durham** & Melville (2) p. 253.

†Pygurus (Pygurus) jurensis, Kongiel p. 73 pl. vii figs. 1-3.

†Pyrina pygaea, Aliev pp. 1171-3 fig. 4.

†Rhabdocidaris sp. cf. zitteli, Eocene British Somaliland, **Kier** (4) p. 844 pl. oiii figs. 1, 2; R. vidali, **Bata**ller p. 32 pl. vi figs. 2-4.

Rotula deciesdigitata and orbiculus, **Tortonese** (1) pp. 1-4.

Rotulina, **Durham & Melville** (2) p. 259, diagnosed.

†Rhynobrissus cuneus sp. nov. North Carolina, U.S., Cooke (2) pp. 10-12 pl. i.

†Salenia geometrica, Chavan & Cailleux p. 194 fig. 2226.

†Schizaster (Schizaster) pp. 880-881 with S. africanus pp. 881-882 pl. ev fig. 11 Kier (4); S. (Paraster) p. 882 with S. (Paraster) duroensis Paleocene British Somaliland pp. 885-886 pl. evi figs. 11-14. hunti Paleocene British Somaliland pp. 882-883 pl. evi figs. 5-7 karkarensis Eocene British Somaliland pp. 883-884 pl. evi figs. 1-3 spp. nov., S. (P.) beloutchistanensis pp. 884-885 pl. cvi figs. 8-10, S. (P.) sp. cf. meslei p. 883 pl. evi fig. 4, Id: S. miyazakiensis sp. nov. Miocene Japan, Morishita (1) pp. 197-199 pl. iii figs. 1a-d, 2a-d, pl. iv figs. la-d; S. vidali, Bataller p. 32 pl. vi figs. 5, 5a.

†Scutella montagnai sp. nov. Oligo-Miocene Italy, Mirigliano pp. 9-12 textfig. 1 pl. i figs. 1, 2; S. subrotunda, Chavan & Cailleux p. 199 fig. 2247.

Scutellina, **Durham & Melville (2)** p. 259, diagnosed.

†Sismondia polymorpha, Cooke (1) pp. 362–363 pl. exix figs. 9–13.

Spatangoida, **Durham & Melville (2)** p. 261, diagnosed.

†Spatangoides striatoradiatus, Chavan & Cailleux p. 204 fig. 2265.

Sperosoma biseriatum, Baranova p. 210.

†Stomechinus bigranularis, Chavan & Cailleux p. 195 fig. 2230a-c; S. gyratus pp. 68-69 pl. v figs. 2-4, S. cf. semiplacenta p. 69 pl. v fig. 1 pl. iv fig. 7, Kongiel.

Strongylocentrotus djakonovi sp. novpp. 223–225 fig. 14. S. polyacanthus apicimagis subsp. nov., Bering Sea, Baranova, S. droebachiensis pp. 212–214 fig. 9, echinoides pp. 216–219 fig. 11, polyacanthus pp. 219–221 fig. 12, pulchellus pp. 211–212 fig. 8, sachalinicus pp. 214–216 fig. 10, Id.; S. nudus, Chang & Woo pp. 131–134 figs. 4–7 pl. i fig. 2 pl. ii fig. 2.

Stylocidaris affinis, Tortonese (3) p. 192

†Temnechinus turbinatus, Engel in Burck p. 36 fig. 2.

Temnopleuroida, Durham & Melville (2) p. 255, diagnosed.

†Tetragramma pomeraniae sp. nov. Jurassic Pomerania, Kongiel pp. 65-67 pl. iv figs. 1-3.

†Thylechinus vidali, Bataller pp. 29-30 pl. v figs. 1, 2.

†Toxaster retusus, Chavan & Cailleux p. 206 fig. 2273.

†Toxaster (Pliotoxaster) collegnei, Aliev pp. 1173-1174 fig. 5.

†*Triplacidia vidali*, Bataller p. 33 pl. vii figs. 1, 1a, 1b.

†Trochalosoma vidali, Bataller p. 29 pl. iv figs. 10-13.

†Trochotiara mamillanum, Kongiel pp. 63-65 pl. iii figs. 4-7.

Tromicosoma tenue, Baranova pp. 210–211.

Urechinus loveni p. 231, narcsianus pp. 229-230, Baranova.

†Vaquerosella coreyi pp. 630-631 pl. lxxii fig. 11, V. sp. indet., Miocene Oregon, U.S., p. 630 pl. lxxii fig. 2, Durham.

†SOMASTEROIDEA

[Vacant]

ASTEROIDEA

Acanthaster planci, Hayashi (1) pp. 169–171.

Aleutiaster schefferi, Baranova p. 166. Allostichaster capensis, Madsen (1) p. 44.

Anasterias antarctica pp. 38–40 pl. v figs. 1, 2, minuta p. 40, varium pp. 40–42 pl. v figs. 3–9, Madsen (1).

Anseropoda placenta, Zavodnik pp. 121-122 fig.

Anteliaster microgenys nannodes, Baranova p. 176.

†Archasterina gen. nov. p. 111, family Palasterinidae, for type A. cornuta sp. nov. Devonian Germany, pp. 111-115 fig. 24 pl. xxxvi figs. 1-5, Lehmann.

†Asteriacites lumbricalis, Jörg pp. 34-35.

Asterias microdiscus p. 178, rathbuni p. 179, Baranova; A. rubens, Gruner in Stresemann p. 454 fig. 452; A. forbesi, Guthrie & Anderson p. 489 fig. 16.2; A. rubens, Lehmann pl. xlvii fig. 3. Asterias (Stolasterias) volsatella, Lehmann pl. xxxii fig. 4.

Asterina coronata forma japonica nov. S. Japan; Hayashi (1) pp. 119–120 pl. xi figs. 5, 6, 7, pl. xiii fig. 7. A. batheri p. 119 pl. xiii figs. 5, 6, exigua p. 121, pectinifera pp. 121–122, Id; A. stellifer, Collignon pl. i fig. 2.

Asterodon singularis, Madsen (1) p. 20.

Astrocles actinodetus, Baranova p. 173.

Astropecten marginatus, Bernasconi (1) pp. 67-68 pl. ii figs. 3, 4; A. michaelseni, Collignon pl. i fig. 1; A. aurantiacus, Cumano cover picture; A. irregularis, Gruner in Stresemann p. 454 fig. 454/1; A. ludwigi, referred to Ctenopleura, Hayashi (2) p. 42.

Astrostole platei, Madsen (1) p. 37.

†Baliactis p. 81, with B. scutatus pp. 85–86 pl. xxvi figs. 3–5, tuberatus pp. 83–84 pl. xxvi figs. 1, 2 spp. nov. Devonian Germany, Lehmann. B. devonicus pp. 81–83 fig. 22a pl. xxv figs. 1–3, B. sp. p. 86 pl. xxvii figs. 1–3, Id.

Benthopecten claviger pp. 158-159 claviger occidentalis p. 159, Baranova.

Brisingella pannychia, Baranova p. 173.

Ceramaster arcticus, clarki p. 162, japonicus, patagonicus p. 161, stellatus pp. 161–162, Baranova; C. granularis, Blacker pp. 18–21, 45 fig. 27a, b.

Cladaster validus, Baranova pp. 162–163.

Coscinasterias tenuispina, Tortonese (3) p. 191.

Cosmasterias lurida, Madsen (1) pp. 42-43 pl. vi figs. 1, 2.

Crossaster borealis p. 165, papposus p. 166, Baranova; C. papposus pp. 186–187, papposus forma japonicus p. 187, Hayashi (1).

Ctenodiscus crispatus, Baranova p. 153.

Ctenopleura fisheri sp. nov. Japan, Hayashi pp. 37-42 text-figs. 1, 2, pl. i figs. 1-3. C. ludwigi pp. 42-44 text-fig. 3 pl. i figs. 4, 5, Id.

Cycethra verrucosa, Madsen (1) p. 21.

Diplopteraster multipes, Baranova p. 167; D. multipes, Hayashi (1) pp. 199– 200 pl. xii fig. 1, pl. xiii fig. 1. Dipsacaster anoplus pp. 156-157,

borealis p. 157. Baranova.

Echinaster antonioensis, Bernasconi (1) pp. 72-73 pl. vi figs. 1, 2; E. luzonicus, Hayashi (1) pp. 123-124; E. sepositus, Tortonese (3) p. 190.

†Echinasterella p. 115, with E. sladeni pp. 116-118 pls. xxxvii-xxxix, Lehmann.

†Encrinaster schmidi, Grabert & Grabert pp. 190-197 fig. 1 pls. xii, xiii.

Enoplopatiria marginata, Bernasconi (1) pp. 70–72 pl. iv fig. 4, pl. v fig. 2 pl. vii fig. 1.

†Eostella gen. nov. p. 146 (incertae sedis) for type E. hunsrückiana sp. nov. Devonian Germany pp. 146-147 fig. 30 pl. 1 figs. 4-6 Lehmann.

Eremicaster pacificus pp. 154-155, tenebrarius p. 154, Baranova.

Evasterias echinosoma p. 178, retifera pp. 177–178, troschelii p. 178, Baranova.

† Forbesiaster gaensis sp. nov. Cretaceous Egypt, Jux pp. 187–189 pl. xxi figs. 1, 2.

Freyella hexactis sp. nov. Bering Sea, Baranova pp. 174-175 fig. 1.

Ganeria falklandica, Madsen (1) p. 22.

Gephyreaster swifti, Baranova p. 159.

†Helianthaster pp. 89-91 with H. rhenanus microdiscus var. nov. Devonian Germany p. 93 pl. xxxi figs. 1, 2, H. rhenanus pp. 91-93 pl. xxx figs. 1-4, Lehmann.

Henricia pp. 125-128, with H. exigua pp. 138-140 figs. 19-21 pl. vii figs. 5, 6, irregularis pp. 164-166 figs. 60-63 pl. iv figs. 1, 2, kinkasani pp. 144-146 figs. 28-31 pl. vii figs. 7, 8, pachyderma pp. 160-162 figs. 53-55 pl. vii figs. 1, 2, pacifica figs. 40-43 pl. ix figs. 7-10, regularis pp. 137-138 figs. 15-18 pl. x figs. 3, 4, reniossa pp. 132-134 figs. 7–10 pl. viii figs. 16, 17, reticulata pp. 162–164 figs. 56–59, pl. ix figs. 1, 2, saghaliensis pp. 142-144 figs. 25-27 pl. vii figs. 3, 4, spp. nov., H. ohshimai forma acutispina pp. 158–160 figs. 50–52 pl. x figs. 5, 6, reniossa forma tohukuensis pp. 135-137 figs. 11-14 pl. vii figs. 11, 12 formae nov. Japan, Hayashi (1); H. aspera pp. 154-156 figs. 44-46 pl. vii figs. 9, 10, densispina pp. 140–142 figs. 22–24 pl. x figs. 7, 8, leviuscula pp. 128-130 figs. 1-3 pl. viii figs. 14, 15, leviuscula spiculifera pp. 130–132 figs. 4–6 pl. ix figs. 3, 4, nipponica pp. 146–149 figs. 32–35 pl. viii figs. 1, 2, 3, 4, ohshimai pp. 156–158 figs. 47–49 pl. ix figs, 5, 6, tumida pp. 149–152 figs. 36–40, pl. viii figs. 5–13, Id; H. arctica p. 172, aspera pp. 171–172, asthenactis p. 173, beringiana p. 172, derjugini p. 172, longispina aleutica pp. 172–173, orientalis p. 172, spiculifera p. 171, tumida pp. 170–171, Baranova; H. sanguinolenta, Gruner in Stresemann p. 454 fig. 454/2. H. obesa p. 30, studeri p. 31, Madsen (1).

Heliaster canopus p. 36, helianthus pp. 34-35, polybrachius p. 36, Madsen (1); H. helianthus, Schneider p. 75.

Hippasteria colossa, spinosa, Baranova p. 162; H. phrygiana, Blacker pp. 18-21, 45 fig. 27c.

†Hunsrückaster gen. nov. (incertae sedis) with H. peregrinus sp. nov. Devonian Germany pp. 141-144 figs. 27-29 pl. xlix figs. 1-4, Lehmann.

Hymenaster koehleri p. 170, perissonotus p. 170, quadrispinosus pp. 169–170, Baranova; H. pellucidus, Blacker pp. 21–22, 46; H. glaucus, Hayashi (1) pp. 200–202 pl. xii fig. 3, pl. xiii figs, 2, 3; H. pellucides, Lehmann pl. xlvii fig. 4.

†Hystrigaster gen. nov. p. 151 (incertae sedis) for type H. horridus sp. nov. Devonian Germany pp. 152-154 pls. lii-lv, Lehmann.

†Jaekelaster p. 135, with J. petaliformis pp. 135-137 pl. xlvii figs. 1, 2 pl. xlviii fig. 1, Lehmann.

†Kyraster gen. nov. p. 144 (incertae sedis) for type K. inermis sp. nov. Devonian Germany pp. 144-145 pl. 1 figs. 1-3, Lehmann.

Labidiaster radiosus, Madsen (1) pp. 33-34 pl. iv fig. 1.

†Leioactis gen. nov. pp. 86-87, family Taeniactinidae, for type L. hunsrückianus sp. nov. Devonian Germany, p. 87 pl. xxvii figs. 4-6, Lehmann.

Leptasterias alaskensis p. 184, alaskensis asiatica pp. 184–185, aleutica p. 185, arctica pp. 179–180, asteira p. 186, camtschatica p. 185, coei truculenta p. 184, groenlandica pp. 181–182, hylodes p. 180, hylodes reticulata pp. 180–181, leptodoma p. 184, ochotensis p. 181, polaris acervata pp. 182–184, squamu-

lata p. 181, stolacantha p. 186, Baranova;
L. polaris, Tortonese (2) pp. 1-5, pls. i,
ii.

Leptychaster anomalus p. 156, arcticus f. microplax p. 155, pacificus pp. 155–156, propinquus p. 156, Baranova.

Lethasterias nanimensis chelifera, Baranova p. 177.

Linckia guildingii, Bernasconi (1) pp. 68-70 pl. v figs. 1, 3.

Lophaster furcilliger vaxator, Baranova p. 163; L. furcifer, Blacker pp. 21–22, 45–46 fig. 28; L. stellans, Madsen (1) pp. 31–32.

Luidia magellanica, Madsen (1) pp. 16-18 pl. i figs. 1, 2; L. sarsi, Tortonese (3) p. 190.

Luidiaster dawsoni, Baranova pp. 157-158.

Marthasterias glacialis, Millott (2) figs. 7, 8, 11.

Mediaster trinidadensis sp. nov. Trinidad Island, off Brazil, Bernasconi (2) pp. 33-34 fig.

†Medusaster p. 94 with M. rhenanus pp. 94-99 fig. 23 pl. xxxii figs. 1-10, Lehmann.

Meyenaster gelatinosus, Madsen (1) p. 36.

†Metopaster parkinsoni, Chavan & Cailleux p. 192 fig. 2218.

Mithrodia clavigera, Hayashi (1) pp. 171-173.

Myxoderma sacculatum, Baranova p. 175.

Nearchaster pedicellaris, variabilis, Baranova p. 158.

Nectria ocellifera, Lehmann pl. xlii figs. 4, 5.

Odontaster mediterraneus, Gautier-Michaz pp. 80-81; O. penicillatus, Madsen (1) pp. 19-20.

†Palaeactis gen. nov. pp. 87-88, family Taeniactinidae, for type P. lanceolatus sp. nov. Devonian Germany pp. 88-89 fig. 226 pl. xxviii figs. 1-4, Lehmann.

†Palaeosolaster pp. 118-119, with P. gregoryi pp. 120-121 pl. xl figs. 1, 2, Lehmann.

† Palaeostella pp. 121–123, with P^* solida pp. 123–126 pl. xli figs. 1–4 pl. xlii figs. 1–5, Lehmann.

†Palasterina p. 100 with P. marginata pp. 108–109 pl. xxxiv figs. 4–6, maucheri pp. 107–108 pl. xxxiv figs. 1–3, taenibrachiata p. 109 pl. xxxv figs. 1–3 tilmanni pp. 104–106 pl. xxix fig. 2 pl. xxxiii figs. 1, 2, spp. nov. Devonian Germany, Lehmann. P. follmanni pp. 100–103 pl. xxiii figs. 3–5, Id.

†Palasteriscus p. 78, with P. devonicus pp. 78–80 pl. xxiv figs. 1, 2, Lehmann.

†Paleaster sp., Ladd p. 36 pl. iii fig. 1.

Patiria chilensis pp. 23-24 pl. iv figs. 2, 3, obesa p. 24 pl. iii figs. 1-5, Madsen (1).

Patiriella calcarata p. 25, fimbriata p. 26, Madsen (1).

Pedicellaster magister, Baranova p. 176.

Peribolaster biserialis, Baranova pp. 166-167.

Porania antarctica, Madsen (1) pp. 27-28.

Poraniopsis p. 166 with P. inflata pp. 167-169 pl. iv figs. 9, 10, Hayashi (1); P. echinaster, Madsen (1) pp. 29-30 pl. ii figs. 1-4.

†Promopalaeaster p. 905 with P. pricei sp. nov. Ordovician Pennsylvania, U.S., pp. 905-906 pl. cviii figs. 3, 4, Cramer.

†Protasteracanthion emended, pp. 147–148, with P. primus pp. 148–151 fig. 31 pl. li figs. 1–4, Lehmann.

Pseudarchaster dissonus pp. 160-161, parellii pp. 159-160, Baranova; P. parelii, Blacker pp. 18-21, 45.

Psilaster pectinatus, Baranova p. 157

Pteraster, key to Japanese spp. pp. 188–189, P. uragaensis sp. nov. pp. 194–195 pl. xi. figs. 3, 4, tesselatus glomus subsp. nov. pp. 193–194 pl. xi figs. 1, 2, Japan, Hayashi (1); P. japonicus pp. 196–197, jordani pp. 195–196 pl. xi fig. 8, pl. xii fig. 2, multiporus pp. 190–191, obscurus pp. 198–199 pl. xii figs. 4–6, pl. xiii fig. 4, obesus pp. 189–190, tesselatus pp. 191–193, Id; P. marsippus p. 168, militaris p. 167, obscurus p. 169, pulvillus p. 168, temnochiton pp. 168–169, tesselatus p. 168, Baranova.

Pycnopodia helianthoides, Baranova pp. 176-177.

†Recurvaster lorioli sp. nov. Cretaceous Egypt, Jux pp. 189-190 fig. 1 pl. xxi fig. 3.

†Schlüteraster gen. nov. pp. 137-138, family Compsasteridae, for type Astropecten schlüteri Stuertz, Lehmann. S. schlüteri pp. 138-141 fig. 26 pl. xlviii figs. 2, 3, Id.

Solaster dawsoni p. 164, endeca pp. 163–164, hypothrissus p. 165, pexillatus p. 164, stimpsoni p. 164, Baranova; S. papposus, Gruner in Stresemann p. 455 fig. 455/1; Solaster key to Japanese spp. pp. 174–175, with S. borealis pp. 183–184, dawsoni pp. 178–180, dawsoni var. intermedius pp. 180–181, endeca pp. 175–177, paxillatus pp. 181–182, stimpsoni pp. 177–178 Uchidai pp. 184–185, Hayashi (1).

Stephanasterias albula, Baranova p.

Stichaster striatus, Madsen (1) pp. 37-38.

†Trichasteropsis weissmanni, Heller pp. 45-47 fig. 1.

†Urasterella pp. 127-128, with U. verruculosa sp. nov. Devonian Germany pp. 131-134 fig. 25 pl. xlv figs. 1-3 pl. xlvi figs. 1-4, Lehmann. U. asperula pp. 128-131 pl. xliii figs. 1-4 pl. xliv figs. 1-6, Id.

Zoroaster actinocles, ophiurus, Baranova p. 175.

OPHIUROIDEA

Acrocnida semisquamata, Cherbonnier (3) p. 167; A. brachiata, Gruner in Stresemann p. 456 fig. 456/3.

Amphiodia craterodmeta p. 197, euryaspis pp. 197–198, periercta p. 197, Baranova.

Amphipholis squamata, Balinsky pp. 10–11; A. nudipora, Cherbonnier (3) p. 167; A. squamata, Gruner in Stresemann p. 456 fig. 456/2.

Amphioplus ailsaclarki sp. nov., Sierra Leone, Cherbonnier (3) pp. 167– 170 figs. 2, 3. A. cincta, Id. p. 167; A. integer, Balinsky p. 11; A. macraspis, Baranova pp. 196–197.

Amphiura inhacensis pp. 5-7 text-fig. 2 pl. iv figs. 13, 14, kalki pp. 3-5 text-fig. 1 pl. i figs. 1, 2 spp. nov. Portuguese E. Africa, Balinsky; A. angularis p. 5, Id; A. apicula sp. nov.,

Mediterranean, Cherbonnier (1) pp. 200-204 figs. 1-3; A. beringiana pp. 199-200 fig. 3, carchara, leptodoma, psilopora p. 198, sundevalli pp. 198-199, Baranova; A. sculpta, Cherbonnier (3) p.167.

Amphylicus androphorus, Balinsky p. 11.

Anthophiura axiologa, Baranova pp. 201–202 fig. 5.

Asteronyx loveni, Baranova p. 186.

Astrochele laevis, Baranova pp. 186–187.

Astrocladus coniferus, Baranova p. 187. Astrophiura chariplax, Baranova pp. 200-201 fig. 4.

†Bundenbachia p. 43 with B. beneckei pp. 43-46 fig. 12 pl. x figs. 1-5 pl. xi figs. 1-4, Lehmann.

†Cheiropteraster p. 35 with C. giganteus pp. 35-38 figs. 9, 10 pl. vii figs. 1, 2, Lehmann.

†Encrinaster pp. 28-29, with E. laevidiscus sp. nov. Devonian Germany, p. 31 pl. v figs. 3-5, roemeri pp. 29-31 figs. 5-8 pl. v figs. 1, 2, Lehmann.

†Eospondylus p. 57 with E. primigenius compactus var. nov. Devonian Germany pp. 60-61 pl. xvi figs. 1-3, E. primigenius pp. 57-60 fig. 16 pl. xv figs. 1-6, Lehmann.

†Erinaceaster gen. nov. family Pradesuridae, p. 20 for type E. tenuispinosus sp. nov. Devonian Germany pp. 20-21, pl. i figs. 1-4, Lehmann; E. giganteus sp. nov. Devonian Germany pp. 22-23 fig. 2 pl. ii fig. 1 pl. iii fig. 1, spinosissimus pp. 21-22 pl. iv figs. 2, 3, Id.

†Euzonosoma pp. 24-25 with E. tischbeinianum pp. 25-28 figs. 3, 4 pl. iv figs. 1, 4-6, Lehmann.

†Furcaster pp. 63-65, with F. decheni pp. 69-72 figs. 18, 20b pl. xix figs. 1-3 pl. xx figs. 1, 2, palaeozoicus pp. 65-69 figs. 17, 20a pl. xviii figs. 1-6, zitteli pp. 72-74 figs. 19, 20c pl. xxi figs. 1-2 pl. xxii figs. 1-3, Lehmann.

†Geocoma elegans, Chavan & Cailleux p. 192 fig. 2217.

Gorgonocephalus caryi, Baranova pp. 187–188; G. arcticus pp. 22–26, 46–47 fig. 29d, caput-medusae pp. 22–26, 46–47 fig. 29a, eucnemis pp. 22–26, 46–47 fig. 29e, lamarcki pp. 22–26, 46–47 fig. 29b, Blacker.

†Hymenosoma gen. nov. p. 31, family Euzonosomatidae, for type H. opitzi sp. nov. Devonian Germany pp. 32-33 pl. vi figs. 1, 2, Lehmann.

†Kentrospondylus gen. nov. p. 61, family Hallasteridae, for type K. decadactylus sp. nov. Devonian Germany, pp. 61-62 fig. 15 pl. xvii, Lehmann.

† Loriolaster p. 38 with L. gracilis sp. nov. Devonian Germany pp. 41-42 pl. ix figs. 3, 4, L. mirabilis pp. 39-41 fig. 11 pl. viii figs. 1-3 pl. ix figs. 1, 2, 5, Lehmann.

Macrophiothrix mossambica sp. nov. Portuguese E. Africa. Balinsky pp. 18-20 text-fig. 7 pl. iii figs. 11, 12. M. aspidota p. 18, hirsuta pp. 17-18, longipeda p. 17, Id.

†Mastigophiura gen. nov. pp. 49-50, family Protasteridae, for type M. grandis sp. nov. Devonian Germany pp. 50-52 fig. 14 pl. xiii figs. 1, 2, Lehmann.

†Miospondylus p. 53 with M. rhenanus pp. 54-57 pl. xiv figs. 3, 4, Lehmann.

Ophiacantha adiaphora p. 191, O. (?) atopostoma p. 190, bathybia p. 192, bidentata p. 191, enneactis pp. 192-193, macrarthra p. 190, nutrix p. 193 fig. 2, rhachophora p. 191, trachybactra pp. 191-192, Baranova.

Ophiacanthella aconotophora, Baranova p. 194.

Ophiactis delagoae sp. nov. Portuguese E. Africa, Balinsky pp. 12-13 text-fig. 5 pl. ii figs. 7, 8; O. carnea pp. 11-12, hemiteles p. 13, lymani, modesta p. 14, parva p. 15, savignyi p. 14, Id; O. lutkeni, Cherbonnier (3) pp. 166-167.

Ophiocentrus dilatatus, Balinsky p. 7.

Ophiocoma erinaceus pp. 25-26, insularia p. 26, parva p. 27, pica p. 26, scolopendrina p. 25, valenciae p. 27, Balinsky.

Ophiocomella clippertoni, Hertlein & Emerson p. 6.

Ophiocten sericeum, Gautier-Michaz

†Ophioglypha graysonensis, Ladd p. 36 pl. iii fig. 3.

Ophiolepis cincta, Balinsky p. 28; O. affinis, Cherbonnier (3) p. 170.

Ophioleuce oxycraspedon, Baranova pp. 208-209 fig. 7.

Ophiolimna bairdi p. 189, papillata pp. 188-189, Baranova.

Ophiomastix notabilis p. 27, venosa pp. 27-28, Balinsky.

Ophionephthys africana sp. nov. Portuguese E. Africa, Balinsky pp. 7-9 text-fig. 3 pl. i figs. 3, 4.

Ophionereis australis, porrecta, vivipara, Balinsky p. 24.

Ophiopenia disacantha p. 208, tetracantha pp. 207-208, vicina p. 208, Baranova.

Ophiopezella decorata, Balinsky p. 28.

Ophiopholis aculeata, Baranova pp. 195-196, Gruner in Stresemann p. 456 fig. 456/1, Guthrie & Anderson fig. 16.7.

Ophiophragmus sacensis sp. nov. Portuguese E. Africa, Balinsky pp. 9-10 text-fig. 4, pl. ii figs. 5, 6.

Ophiophthalmus cataleimmoidus p. 190, normani p. 189, Baranova

Ophiopleura borealis, Blacker pp. 26, 47 fig. 30a.

Ophiopteron atlanticum, Cherbonnier (3) pp. 163–166 fig. 1.

Ophioschiza monacantha, Baranova p. 188.

Ophiosemnotes brevispina, diaphora, pachybactra, paucispina p. 195, tylota p. 194, Baranova.

Ophiothela beauforti pp. 22-24 pl. iv fig. 16, dividua p. 22, Balinsky.

Ophiothrix echinotecta sp. nov. Portuguese E. Africa, Balinsky pp. 16-17 text-fig. 6, pl. iii figs. 9. 10; O. spiculata, Baranova p. 188.

Ophiotrichoides propinqua, Balinsky pp. 21-22.

Ophiura bathybia pp. 205-206, cryptolepis p. 207, flagellata pp. 206–207, irrorata p. 207, leptoctenia p. 205, maculata p. 204, quadrispina p. 206, sarsi p. 205, Baranova; O. africana, Cherbonnier (3) pp. 170-171; O. affinis p. 457 fig. 457/3, albida p. 457 fig. 457/2, texturata p. 457 figs. 455/2, 457/1, Gruner in Stresemann.

†Ophiurina pp. 74-75 with O. lymani pp. 75-77 fig. 21 pl. xxiii figs. 1-3, Lehmann.

† Palaeophiomyxa pp. 46-47 with P. grandis pp. 47-49 fig. 13 pl. xii figs. 1-5, Lehmann.

†Palaeophiura simplex, Lehmann pp. 52–53 pl. xiv figs. 1, 2.

Placophiothrix foveolata p. 20 pl. iv fig. 15, proteus, trilineata p. 21, Balinsky.

26

Stegophiura nodosa, **Baranova** pp. 202–204 fig. 6.

†Taeniaster p. 906 with T. spinosus pp. 906-907 pl. cviii figs. 1, 2 (T. maximus Willard a synonym), Cramer.

CRINOIDEA

† Acanthocrinus rex. Struve fig. 1.

†Allagecrinus sculptus sp. nov. Carboniferous New Mexico, Strimple & Koenig pp. 1229–1230 fig. 2₁₋₁₂.

†Amphipsalidocrinus, Strimple & Koenig pp. 1239-1240.

Antedon bifida (Pennant) 1777 a senior subjective synonym of Antedon gorgonia Fréminville, 1811 and type species of Antedon Fréminville, 1811, Direction 73 p. 209; A. rosaceus, Guthrie & Anderson fig. 16.15A, B; Antedon, Struve fig. 14.

†Apiocrinus parkinsoni, Chavan & Cailleux p. 191 fig. 2214.

Bathycrinus gracilis Thomson 1872, placed on the Official List of Specific Names in Zoology, Direction 73 pp. 198, 208; B. complanatus, Baranova p. 153.

Bathymetra brevicirra, Baranova p. 153.

†Botryocrinus ? sp., Sieverts-Doreck (1) p. 66.

†Bourgueticrinus includes Mesocrinus Carpenter, 1881 and Metapiocrinus Jackel, 1918, Rasmussen p. 554.

†Camarocrinus mariannae sp. nov., Silurian Kazakhstan, U.S.S.R., Yakovlev (4) p. 29 pl. i figs. 1, 2 (see also Scyphocrinus).

†Cromyocrinus simplex, Yakovlev (6) p. 12 figs. 3a, 6.

†Crotalocrinites baschkiricus sp. nov. Silurian or Devonian U.S.S.R., Yakov-lev (3) p. 15 pl. i figs. 1, 2.

†Ctenocrinus sp. nov. Devonian Germany, Sieverts-Doreck (1) pp. 65-66.

†Cupressocrinus crassus, Chavan & Cailleux p. 189 fig. 2205; Cupressocrinus, Firtion pp. 163–164 fig. 9 pl. x figs. 19, 20; C. abbreviatus, Struve fig. 13.

† Dactylocrinus p. 55 with D. oligoptilus pp. 55-56 pl. x figs. 1, 2, Yakovlev (2). †Decacrinidae fam. nov, Eltuischeva p. 220.

†Decacrinus gen. nov. p. 221, family Decacrinidae, for type D. pennatus sp. nov. Devonian Kazakhstan, U.S.S.R. pp. 221-222 fig. 1 pl. i figs. 1-4, Eltuishcheva. D. orientalis sp. nov. Devonian Amur, U.S.S.R. pp. 222-223 fig. 2 pl. i figs. 5-7, Id.

†Diamenocrinus sp. cf. opitzi, Sieverts-Doreck pp. 63-65 figs. 1a-b.

† Dichostreblocrinus pyriformis sp. nov. Carboniferous New Mexico, U.S., Strimple & Koenig pp. 1243-1244 fig. 4_{10-15} .

†Encrinus liliiformis, Chavan & Cailleux p. 190 fig. 2213; E. granulosus, Szörényi p. 130.

†Eucalyptocrinus rosaceus, Chavan & Cailleux p. 189 fig. 2207.

Florometra asperrima, Baranova p. 152.

†Glenotremites janeti referable to Orthogonocrinus (Roveacrinidae), Rasmussen p. 555.

Heliometra glacialis, Blacker pp. 16-18, 45 fig. 26a.

Holopus rangii d'Orbigny 1837, placed on the Official List of Specific Names in Zoology, **Direction 73** pp. 198, 208.

†Indocrinus crassus fig. 2Γ, elegans fig. 2π and piszowi fig. 2Β, Yakovlev (6).

†Isocrinus agassizii and carinatus, Rasmussen pp. 553-554.

 $\dagger Isselicrinus$ with type I. insculptus, Rasmussen p. 553.

 $\dagger Jimbacrinus$ bostocki, Yakovlev (6) pp. 11–12 figs. 1a, 5.

†Kallimorphocrinus pp. 1230–1233 with K. angulatus. Carboniferous New Mexico, U.S. pp. 1233–1234 figs. 1c, 2₁₃₋₂₈, tintinabulum p. 1235 figs. 1d, 2₃₈₋₃₉, weldenensis pp. 1234–1235 fig. 2₂₈₋₃₅, both Carboniferous Oklahoma, spp. nov. Strimple & Koenig; K. pristinus fig. 1a, puteatus fig. 1b, Id.

†Kuzbassocrinus gen. nov. p. 223, family Decacrinidae, for type K. bystrowi sp. nov. Devonian Kuzbass, U.S.S.R. pp. 223-224 fig. 3 pl. ii figs. 1-2a, Eltuishcheva; K. binidigitatus Devonian Altai, U.S.S.R. pp. 224-225 fig. 4 pl. ii figs. 3-3B, decembotatus Devonian Amur, U.S.S.R. pp. 225-226

fig. 5 pl. i figs. 8-9a, paucicostalus Devonian Pribalkhash, U.S.S.R. pp. 226-227 fig. 6 pl. ii figs 4-5a spp. nov., Id.

†Lampadosocrinus gen. nov. pp. 1244–1245, family Codiacrinidae, for type Dichostreblocrinus minutus Peck, 1936, L. obtusus sp. nov. Carboniferous New Mexico, U.S. pp. 1245–1246 fig. 4_{16-18} , Strimple & Koenig; L. minutus p. 1245 fig. 4_{19-21} , Id.

†Laudonocrinus sp. undet., Strimple p. 369 fig. 1.

†Lipocrinus a synonym of Isselicrinus, Rasmussen p. 553.

†Marsupites testudinarius, Struve fig. 3.

†Mesocrinus Carpenter, 1881 a synonym of Bourgeticrinus, Rasmussen p. 554.

Metacrinus wyvillii Carpenter 1884, placed on the Official List of Specific Names in Zoology, **Direction 73** pp. 198, 208.

†Metapiocrinus Jaekel, 1918 a synonym of Bourgueticrinus, Rasmussen p. 554.

†Millericrinus horridus, Chavan & Cailleux p. 191 fig. 2215.

†Ottowacrinus typus, Sieverts-Doreck (2) figs. 3-4.

†Pagecrinus p. 56 with P. heckeri p. 56 pl. x fig. 3, Yakovlev (2).

†Parabotryocrinus tschudowensis, Yakovlev (2) p. 57 pl. x fig. 4.

†Parapernerocrinus gen. nov. p. 17 family Crotalocrinitidae, for type P. sibiricus sp. nov. Devonian Siberia, U.S.S.R. pp. 16-18 pl. i figs. 4-8 pl. ii figs. 1, 2, Yakovlev (3).

†Passalocrinus triangularis, Strimple & Koenig p. 1246 fig. 4₂₂₋₂₇.

†Pentacrinus tuberculatus, Chavan & Cailleux p. 191 fig. 2216; P. crassus Nielsen, 1913 invalidated by P. crassus Desor, 1845 but referable to Isselicrinus Rovereto, Rasmussen p. 553.

†Plussacrinus gen. nov. family Decacrinidae, für type P. flabellum sp. nov. Ordovician Leningrad Province, Estonia, U.S.S.R. pp. 228–229 fig. 8 pl. iii figs. 2–2B, Eltuishcheva; P. dentatus sp. nov. Ordovician Leningrad Province pp. 229–230 fig. 9 pl. iii figs. 3–3B, Id.

†Podoliocrinus gen. nov. p. 227, family Decacrinidae, for type P. niki-forovae sp. nov. Silurian Podolia, U.S.S.R. pp. 227-228 fig. 7 pl. iii figs. 1-1B, Eltuishcheva.

Poliometra prolixa, Blacker pp. 16-18, 45 fig. 26b.

†Protocrinites yakovlevi sp. nov. Ordovician Leningrad Province & Estonia, Ghekker & Ghekker pp. 274–278 figs. 1-4.

Psathyrometra fragilis, Baranova p. 152.

†Pseudosaccocoma strambergensis, Patrulius pp. 187–191 figs. 1–5.

Retiometra alascana, Baranova p. 152.

†Rhipidocrinus (Rhodocrinus), Firtion pp. 164–166 figs. 10, 11 pl. x figs. 21, 22.

Rhizocrinus lofotensis, correction of entry in Official List from M. Sars 1865 to M. Sars 1868, Direction 72 pp. 166, 170, 171, 176, 177; Direction 73 pp. 198, 208.

†Ristnacrinus, Yakovlev (5) pp. 155-157 figs. 1, 2.

Saccocoma tenella, Struve fig. 4.

†Salagastiana simetrica, Sieverts-Doreck (2) pp. 152-153 figs. 1-2.

†Scyphocrinus mariannae sp. nov. Silurian Kazakhstan, U.S.S.R., Yakovlev (4) p. 29 pl. iii figs. 1-5.

†Sphaerometra Gislén, 1924 possibly identical with Glenotremites Goldfuss 1931, Rasmussen p. 554.

†Syndetocrinus uralicus sp. nov. Silurian U.S.S.R., Yakovlev (3) p. 18 pl. ii figs. 3-5; S. sp. pl. ii fig. 6. Id.

Thalassometra villosa, Baranova p. 153.

† Trophocrinus pp. 1235–1237 with T. bicornis p. 1238 fig. 3_{5-12} , brevis pp. 1238–1239 fig. 3_{13-16} , variabilis pp. 1237–1238 fig. 3_{1-4} spp. nov. Carboniferous Oklahoma, U.S., Strimple & Koenig.

Tropiometra carinata, Tommasi p. 37 figs. 26, 27.

† Tytthocrinus pp. 1240-1241 with T. alamogordoensis sp. nov. Carboniferous New Mexico, U.S. pp. 1241-1243 fig. 4_{1-9} , Strimple & Koenig.

†Ureocrinus bockschii, Yakovlev (6) p. 12 figs. 2a, ő. †Uintacrinus socialis, Chavan & Cailleux p. 190 fig. 2212, Ladd p. 52 pl. ix fig. 1, Miller, Sternberg & Walker pp. 163-166 fig. 1, Struve p. 363 figs. 5, 6, 8, 10, 11, 12.

†MACHAERIDIA

[Vacant.]

†EOCRINOIDEA

Mimocystites bohemicus, Choubert & Termier pp. 138-142 figs. 1-7.

†PARACRINOIDEA

[Vacant.]

†CYSTOIDEA

Echinosphaerites aurantium, Chavan & Cailleux p. 187 fig. 2200, Vokes (1) pl. i figs., 9, 10.

"Eocystites" sp. indet., Vermont, U.S., Shaw p. 788 pl. xeviii fig. 20.

Pleurocystites, Guthrie & Anderson fig. 161A.

Pseudocrinites gordoni, Vokes (1) pl. iv figs. 10, 11.

†CARPOIDEA

Paranacystis petrii, Caster & Eaton pp. 611-614 figs. 1, 2B, C.

†EDRIOASTEROIDEA

[Vacant.]

†BLASTOIDEA

Blastoidocrinus, Guthrie & Anderson fig. 16.1B.

Cryptoblastus submelo sp. nov. Carboniferous Kazakhstan, U.S.S.R., Yakovlev (1) p. 72 pl. x fig. 5.

Nymphaeoblastus kasakhstanensis sp. nov. Carboniferous Kazakhstan, U.S.S.R. Yakovlev (1) p. 71 fig. 5 pl. x fig. 3; N. anossofi p. 71 pl. x fig. 1, miljukovi p. 71 pl. x fig. 2, Id.

Orbitremites coralridgensis pp. 137–139 fig. 1A pl. xiv figs. 8–11, Kentuckyensis pp. 135–137 figs. 1B, 1C, 1E pl. xiii figs. 1–20 spp. nov. Carboniferous Kentucky, U.S. Conkin; O. oppeltiemended, pp. 139–140 fig. 1D pl. xiv figs. 1–7, Id.

Г1957]

Pentremites pp. 1–104 with P. gutschicki p. 68 pl. vii figs. 2, 3 pl. xii figs. 13–21, halli pp. 71–72 pl. ix figs. 2–7 pl. xiii figs. 4, 5 both Carboniferous Kentucky, U.S., malotti p. 48 pl. iii figs. 9, 10 Carboniferous Indiana, U.S. spp. nov., Galloway & Kaska; P. abruptus pp. 54-55 pl. iv figs. 9-11, altus p. 55 pl. iv figs. 19, 20, angularis p. 64 pl. v figs. 25-27, basilaris p. 74 pl. x figs. 9, 10, biconvexus pp. 47-48 pl. iii figs. 5-8, bradleyi p. 74 pl. x fig. 11, broadheadi p. 71 pl. ix fig. 1, burlingtonensis p. 42 pl. ii fig. 4, buttsi p. 58 pl. v figs. 7-9, cervinus p. 66 pl. vi figs. 9, 10, cherokeeus p. 72 pl. ix figs. 8, 9, chesterensis pp. 69-70 pl. viii fig. 1, clavatus p. 60 pl. v figs. 19-21, conoideus pp. 42-43 pl. ii figs. 8-15 pl. xi figs. 1-19 pl. xiii fig. 8, conoideus amplus p. 44 pl. ii figs. 21-26, conoideus obtusus pp. 43–44 pl. ii figs. 19, 20, conoideus perlongus p. 43 pl. ii figs. 16–18, decussatus pp. 41-42 pl. ii figs. 2, 3, elegans pp. 64-65 pl. v figs. 28-30 pl. vi figs. 1-4, elongatus p. 42 pl. ii figs. 5-7, fohsi p. 66 pl. vi figs. 11-13 pl. xiii fig. 6, gemmiformis p. 58 pl. v figs. 5, 6, girtyi pp. 59-60 pl. i fig. 8 pl. v figs. 15, 16 pl. xi figs. 31-42, globosus pp. 74-75 godoni pp. 48-49 pl. iii figs. 11-13 pl. xi figs. 20-30 pl. xiii figs. 9-12, godoni abbreviatus p. 49 pl. iii figs. 14-17, godoni angustus p. 50 pl. iii figs. 23, 24 pl. xiii fig. 1, godoni major pp. 49–50 pl. iii figs. 20–22, godoni pinguis p. 45 pl. iii figs. 1, 2, grandis p. 75, hambachi pp. 65-66 pl. vi figs. 7, 8, hemisphericus p. 68 pl. vii figs. 4, 5, k bi pp. 58-59 pl. v figs. 10-12, laminatus pp. 67-68 pl. vii fig. 1, maccalliei p. 70 pl. viii figs. 2, 3, nodosus p. 65 pl. vi figs. 5, 6, obesus pp. 70-71 pl. viii figs. 4-6, okawensis p. 60 pl. v figs. 17, 18 pl. xi figs. 43-54, ovalis pp. 44-45 pl. ii fig. 29, ovoides p. 44 pl. ii figs. 27, 28, patei p. 57 pl. v figs. 1, 2, perelongatus p. 75, platybasis pp. 50-51 pl. iii figs. 25-29, princetonensis pp. 52-54 pl. iv figs. 5-8, pulchellus pp. 51-52 pl. iv figs. 1-4, pyramidatus p. 57 pl. v figs. 3, 4 pl. xiii fig. 7, pyriformis pp. 56–57 pl. iv figs. 32–37, pl. xiii figs, 2, 3, robustus pp. 66–67 pl. vi figs. 14, 15, rusticus p. 50 pl. iii figs. 18, 19, serratus pp. 73–74 pl. x figs. 7, 8, speciosus p. 59 pl. v fig. 14, spicatus pp. 72–73 pl. ix figs. 10, 11 pl. x figs. 1–4, spicatus porrectus p. 73 pl. x figs. 5, 6, spinosus p. 69 pl. vii figs. 8, springeri pp. 62–64 pl. vii figs. 22–24, sulcatus pp. 68–69 pl. vii figs. 6, 7 pl. xii figs. 1–12, symmetricus p. 55 pl. iv figs. 12–18, tulipaformis p. 67 pl. vi figs. 16, 17, turbinatus p. 59 pl. v fig. 13, tuscumbiae pp. 46–47

pl. iii figs. 3, 4, welleri p. 56 pl. iv figs. 21-31, Id; P. florealis, Chavan & Cailleux p. 188 fig. 2204; probable priority of P. altimarginatus T. H. Clark over biconvexus Ulrich and of divergens Clark over princetonensis Ulrich, Sinclair p. 982.

Pentremitidea leda, Galloway & Kaska pp. 40-41 pl. ii fig. 1.

Schizoblastus librovitchi sp. nov. Carboniferous Urals, U.S.S.R., Yakovlev (1) pp. 71-72 pl. x fig. 4.



VOL. 94.

e 4

